



STUDY GUIDE

MATH AA

SL

www.ib.academy



Mathematics Analysis and Approaches SL Study Guide

Available on www.ib.academy

Author: Nikita Smolnikov, Alex Barancova

Contributing Authors: Laurence Gibbons

Design

Rational

Typesetting



TeX Academy

Special thanks: Robert van den Heuvel

*This work may be shared digitally and in printed form,
but it may not be changed and then redistributed in any form.*

Copyright © 2020, IB Academy

Version: MathAASL.1.0.200507



This work is published under the
Creative Commons BY-NC-ND 4.0 International License.
To view a copy of this license, visit
creativecommons.org/licenses/by-nc-nd/4.0

This work may not be used for commercial purposes other than by IB Academy, or parties directly licenced by IB Academy. If you acquired this guide by paying for it, or if you have received this guide as part of a paid service or product, directly or indirectly, we kindly ask that you contact us immediately.

Daltonlaan 400
3584 BK Utrecht
The Netherlands

www.ib.academy
contact@ib.academy
+31 (0) 30 4300 430

Welcome to the IB.Academy guide for Mathematics Analysis and Approaches SL.

Our Study Guides are put together by our teachers who worked tirelessly with students and schools. The idea is to compile revision material that would be easy-to-follow for IB students worldwide and for school teachers to utilise them for their classrooms. Our approach is straightforward: by adopting a step-by-step perspective, students can easily absorb dense information in a quick and efficient manner. With this format, students will be able to tackle every question swiftly and without any difficulties.

For this guide, we supplement the new topics with relevant sections from our previous Math Studies, SL and HL study resources, and with insights from our years of experience teaching these courses. We illustrate theoretical concepts by working through IB-style questions and break things down using a step-by-step approach. We also include detailed instructions on how to use the TI-Nspire™ to solve problems; most of this is also quite easily transferable to other GDC models.

The best way to apply what you have learned from the guides is with a study partner. We suggest revising with a friend or with a group in order to immediately test the information you gathered from our guides. This will help you not only process the information, but also help you formulate your answers for the exams. Practice makes better and what better way to do it than with your friends!

In order to maintain our Study Guides and to put forth the best possible material, we are in constant collaboration with students and teachers alike. To help us, we ask that you provide feedback and suggestions so that we can modify the contents to be relevant for IB studies. We appreciate any comments and hope that our Study Guides will help you with your revision or in your lessons. For more information on our material or courses, be sure to check our site at www.ib.academy.

IB.Academy Team


If you would like to consider supporting our materials and be recognised for it, send us an email to contact@ib.academy.

PRIOR KNOWLEDGE

Before you start make sure you have a firm grasp of the following. Many marks are lost through errors in these fundamentals.

0.1 Number

Numbers can be grouped in to a number of sets. From the diagram you see that all rational numbers are also real numbers; i.e. \mathbb{Q} is a subset of \mathbb{R} .



Positive integers	$\mathbb{Z}^+ = \{1, 2, 3, \dots\}$
Positive integers and zero	$\mathbb{N} = \{0, 1, 2, 3, \dots\}$
Integers	$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$
Rational numbers	$\mathbb{Q} =$ any number that can be written as the ratio $\frac{p}{q}$ of any two integers, where $q \neq 0$

0.2 Signs

+ and - signs describe positive and negative numbers. Remember they work the opposite way with negative integers. In maths two wrongs do make a right.

$$1 - -1 = 1 + 1 = 2$$

$$-1 \times -1 = 1$$


0.3 Standard form



Standard form is just a way of rewriting any number, sometimes also referred to as 'scientific notation'. This should be in the form $a \times 10^k$, where a is between 1 and 10, and k is an integer.	10	1×10^1
	1000	1×10^3
	3280	3.28×10^3
	4582000	4.582×10^6

0.4 BIDMAS

A handy acronym for remembering the order in which to calculate equations:



Therefore in the following equation

$$\begin{aligned}
 & 4^2 + 5 \times \frac{6}{4} \times (9 - 1) = \\
 \text{B} \quad \longrightarrow & = 4^2 + 5 \times \frac{6}{4} \times (8) = \\
 \text{I} \quad \longrightarrow & = 16 + 5 \times \frac{6}{4} \times 8 = \\
 \text{D/M} \quad \longrightarrow & = 16 + \frac{30}{4} \times 8 = \\
 & = 16 + 7.5 \times 8 = \\
 & = 16 + 60 = \\
 \text{A/S} \quad \longrightarrow & = 76
 \end{aligned}$$

0.5 Solving simultaneous equations

If we have two unknowns, for example x and y , and two equations, then we can solve for x and y simultaneously.

$$\begin{cases} (1) & y = 3x + 1 \\ (2) & 2y = x - 1 \end{cases}$$

There are 3 methods to solve simultaneous equations.

Elimination

Multiply an equation and then subtract it from the other in order to eliminate one of the unknowns.

$$\begin{aligned} 3 \times (2) &\Rightarrow (3) \quad 6y = 3x - 3 \\ (3) - (1) &\Rightarrow 6y - y = 3x - 3x - 3 - 1 \\ &5y = -4 \\ &y = -\frac{4}{5} \end{aligned}$$

Put y in (1) or (2) and solve for x

$$\begin{aligned} -\frac{4}{5} &= 3x + 1 \\ 3x &= -\frac{9}{5} \\ x &= -\frac{9}{15} = -\frac{3}{5} \end{aligned}$$

Substitution

Rearrange and then substitute one in to another.

Substitute (1) into (2)

$$\begin{aligned} 2(3x + 1) &= x - 1 \\ 6x + 2 &= x - 1 \\ 5x &= -3 \\ x &= -\frac{3}{5} \end{aligned}$$


Put x in (1) or (2) and solve for y

$$\begin{aligned} y &= 3\left(-\frac{3}{5}\right) + 1 \\ y &= -\frac{4}{5} \end{aligned}$$

Graph

Graph both lines on your GDC. Where they intersect will be the solution to the equation.

Note that this method is also great when you have to solve more complex equations.



0.6 Geometry

These are given in the data booklet

Area of parallelogram	$A = b \times h$
Area of a triangle	$A = \frac{1}{2}(b \times h)$
Area of a trapezoid	$A = \frac{1}{2}(a + b)h$
Area of a circle	$A = \pi r^2$
Circumference of a circle	$C = 2\pi r$
Volume of a cuboid (rectangular prism)	$V = l \times w \times h$
Volume of a cylinder	$V = \pi r^2 h$
Volume of a prism	$V = Ah$
Area of the curved surface of a cylinder	$A = 2\pi rh$

ALGEBRA

Table of contents & cheatsheet

1.1. Sequences

10

Arithmetic: +/− common difference

$$u_n = n^{\text{th}} \text{ term} = u_1 + (n-1)d$$

$$S_n = \text{sum of } n \text{ terms} = \frac{n}{2}(2u_1 + (n-1)d)$$

with $u_1 = a = 1^{\text{st}}$ term, $d =$ common difference.

Geometric: \times/\div common ratio

$$u_n = n^{\text{th}} \text{ term} = u_1 \cdot r^{n-1}$$

$$S_n = \text{sum of } n \text{ terms} = \frac{u_1(1-r^n)}{(1-r)}$$

$$S_\infty = \text{sum to infinity} = \frac{u_1}{1-r}, \text{ when } -1 < r < 1$$

with $u_1 = a = 1^{\text{st}}$ term, $r =$ common ratio.

Sigma notation

A shorthand to show the sum of a number of terms in a sequence.

$$\sum_{n=1}^{10} 3n-1$$

Last value of n

← Formula

First value of n

e.g.

$$\sum_{n=1}^{10} 3n-1 = \underbrace{(3 \cdot 1) - 1}_{n=1} + \underbrace{(3 \cdot 2) - 1}_{n=2} + \dots + \underbrace{(3 \cdot 10) - 1}_{n=10} = 155$$

1.2. Exponents and logarithms

14

Exponents

$$x^1 = x$$

$$x^m \cdot x^n = x^{m+n}$$

$$(x^m)^n = x^{m \cdot n}$$

$$x^{-1} = \frac{1}{x}$$

$$x^{\frac{1}{2}} = \sqrt{x}$$

$$\sqrt{x}y = \sqrt{x} \cdot \sqrt{y}$$

$$x^{\frac{m}{n}} = \sqrt[n]{x^m}$$

$$x^0 = 1$$

$$\frac{x^m}{x^n} = x^{m-n}$$

$$(x \cdot y)^n = x^n \cdot y^n$$

$$x^{-n} = \frac{1}{x^n}$$

$$\sqrt{x} \cdot \sqrt{x} = x$$

$$x^{\frac{1}{n}} = \sqrt[n]{x}$$

$$x^{-\frac{m}{n}} = \frac{1}{\sqrt[n]{x^m}}$$

Logarithms

$$\log_a a^x = x$$

$$a^{\log_a b} = b$$

Let $a^x = b$, isolate x from the exponent: $\log_a a^x = x = \log_a b$

Let $\log_a x = b$, isolate x from the logarithm: $a^{\log_a x} = x = a^b$

Laws of logarithms

I: $\log_c a + \log_c b = \log_c (a \cdot b)$

II: $\log_c a - \log_c b = \log_c \left(\frac{a}{b}\right)$

III: $n \log_c a = \log_c (a^n)$

IV: $\log_b a = \frac{\log_c a}{\log_c b}$

1.3. Binomial Expansion


16

In a expansion of a binomial in the form $(a+b)^n$. Each term can be described as ${}^n C_r a^{n-r} b^r$, where ${}^n C_r$ is the coefficient.

The full expansion can be written thus

$$(a+b)^n = {}^n C_0 a^n + {}^n C_1 a^{n-1} b + {}^n C_2 a^{n-2} b^2 + \dots + {}^n C_{n-1} a b^{n-1} + {}^n C_n b^n$$

Find the coefficient using either pascals triangle



Or the nCr function on your calculator

1.1 Sequences

1.1.1 Arithmetic sequence



Arithmetic sequence the next term is the previous number + the common difference (d).

e.g. $2, 4, 6, 8, 10, \dots$ $d = +2$ and $2, -3, -8, -13, \dots$ $d = -5$

To find the common difference d , subtract two consecutive terms of an arithmetic sequence from the term that follows it, i.e. $u_{(n+1)} - u_n$.

DB 1.2 Use the following equations to calculate the n^{th} term or the sum of n terms.

$$u_n = u_1 + (n-1)d \qquad S_n = \frac{n}{2}(2u_1 + (n-1)d)$$

with

$$u_1 = a = 1^{\text{st}} \text{ term} \qquad d = \text{common difference}$$

Often the IB requires you to first find the 1^{st} term and/or common difference.

Finding the first term u_1 and the common difference d from other terms.

In an arithmetic sequence $u_{10} = 37$ and $u_{22} = 1$. Find the common difference and the first term.

- | | | |
|----|--|--|
| 1. | Put numbers in to n^{th} term formula | $37 = u_1 + 9d$ $1 = u_1 + 21d$ |
| 2. | Equate formulas to find d | $21d - 1 = 9d - 37$ $12d = -36$ $d = -3$ |
| 3. | Use d to find u_1 | $1 - 21 \cdot (-3) = u_1$ $u_1 = 64$ |

1.1.2 Geometric sequence



Geometric sequence the next term is the previous number multiplied by the common ratio (r).

To find the common ratio, divide any term of an arithmetic sequence by the term that precedes it, i.e. $\frac{\text{second term } (u_2)}{\text{first term } (u_1)}$ e.g. $2, 4, 8, 16, 32, \dots$ $r = 2$

and $25, 5, 1, 0.2, \dots$ $r = \frac{1}{5}$

Use the following equations to calculate the n^{th} term, the sum of n terms or the sum to infinity when $-1 < r < 1$.

DB 1.3 & 1.8

$$\begin{array}{lll}
 u_n = n^{\text{th}} \text{ term} & S_n = \text{sum of } n \text{ terms} & S_\infty = \text{sum to infinity} \\
 = u_1 \cdot r^{n-1} & = \frac{u_1(1-r^n)}{(1-r)} & = \frac{u_1}{1-r}
 \end{array}$$

again with

$$u_1 = a = 1^{\text{st}} \text{ term} \qquad r = \text{common ratio}$$

Similar to questions on Arithmetic sequences, you are often required to find the 1st term and/or common ratio first.

1.1.3 Sigma notation

Sigma notation is a way to represent the summation of any sequence — this means that it can be used for both arithmetic or geometric series. The notation shows you the formula that generates terms of a sequence and the upper and lower limits of the terms that you want to add up in this sequence.



$$\begin{array}{l}
 \downarrow \text{Last value of } n \\
 \sum_{n=1}^{10} 3n - 1 \leftarrow \text{Formula} \\
 \uparrow \text{First value of } n
 \end{array}$$

$$\text{e.g. } \sum_{n=1}^{10} 3n - 1 = \underbrace{(3 \cdot 1) - 1}_{n=1} + \underbrace{(3 \cdot 2) - 1}_{n=2} + \underbrace{(3 \cdot 3) - 1}_{n=3} + \dots + \underbrace{(3 \cdot 10) - 1}_{n=10} = 155$$

Finding the first term u_1 and common ratio r from other terms.

$$\sum_1^5 (\text{Geometric series}) = 3798, \quad \sum_1^{\infty} (\text{Geometric series}) = 4374.$$

Find $\sum_1^7 (\text{Geometric series})$

1. Interpret the question

The sum of the first 5 terms of a geometric sequence is 3798 and the sum to infinity is 4374. Find the sum of the first 7 terms

2. Use formula for sum of n terms

$$3798 = u_1 \frac{1-r^5}{1-r}$$

3. Use formula for sum to infinity

$$4374 = \frac{u_1}{1-r}$$

4. Rearrange **3.** for u_1

$$4374(1-r) = u_1$$

5. Substitute in to **2.**

$$3798 = \frac{4374(1-r)(1-r^5)}{1-r}$$

6. Solve for r

$$\begin{aligned} 3798 &= 4374(1-r^5) \\ \frac{3798}{4374} &= 1-r^5 \\ r^5 &= 1 - \frac{211}{243} \\ \sqrt[5]{r} &= \sqrt[5]{\frac{32}{243}} \\ r &= \frac{2}{3} \end{aligned}$$

7. Use r to find u_1

$$\begin{aligned} u_1 &= 4374 \left(1 - \frac{2}{3}\right) \\ u_1 &= 1458 \end{aligned}$$

8. Find sum of first 7 terms

$$1458 \frac{1 - \left(\frac{2}{3}\right)^7}{1 - \frac{2}{3}} = 4370$$

1.1.4 Compound interest

Sequences can be applied to many real life situations. One of those applications is calculating the interest of a loan or a deposit. Compound interest specifically deals with interest that is applied on top of previously calculated interest. For example, if you make a deposit in a bank and reinvest the interest you will gain even more interest next time. This happens because interest is calculated not just from your initial sum, but also including your re-investments.



$$FV = PV \times \left(1 + \frac{r}{100k}\right)^{kn}$$

DB 1.4

Where:

FV is the future value,

PV is the present value,

n is number of years,

k is the number of compounding periods per year,

$r\%$ is the nominal annual rate of interest

Example.

A deposit of 1000\$ was made in a bank with annual interest of 3% that is compounded quarterly. Calculate the balance in 5 years.

We can use our compound interest equation. Let's identify the known variables.

$$PV = 1000\$$$

$$n = 5$$

$$k = 4$$

$$r = 3\%$$

$$FV = PV \cdot \left(1 + \frac{r}{100k}\right)^{kn}$$

$$FV = 1000 \cdot \left(1 + \frac{3}{100 \cdot 4}\right)^{4 \cdot 5}$$

$$FV = 1160\$$$

1.2 Exponents and logarithms

1.2.1 Laws of exponents

Exponents always follow certain rules. If you are multiplying or dividing, use the following rules to determine what happens with the powers.

Example.

$$\begin{array}{ll}
 x^1 = x & 6^1 = 6 \\
 x^0 = 1 & 7^0 = 1 \\
 x^m \cdot x^n = x^{m+n} & 4^5 \cdot 4^6 = 4^{11} \\
 \frac{x^m}{x^n} = x^{m-n} & \frac{3^5}{3^4} = 3^{5-4} = 3^1 = 3 \\
 (x^m)^n = x^{m \cdot n} & (10^5)^2 = 10^{10} \\
 (x \cdot y)^n = x^n \cdot y^n & (2 \cdot 4)^3 = 2^3 \cdot 4^3 \quad \text{and} \quad (3x)^4 = 3^4 x^4 \\
 x^{-1} = \frac{1}{x} & 5^{-1} = \frac{1}{5} \quad \text{and} \quad \left(\frac{3}{4}\right)^{-1} = \frac{4}{3} \\
 x^{-n} = \frac{1}{x^n} & 3^{-5} = \frac{1}{3^5} = \frac{1}{243}
 \end{array}$$

1.2.2 Fractional exponents

When doing mathematical operations (+, −, × or ÷) with fractions in the exponent you will need the following rules. These are often helpful when writing your answers in simplest terms.

Example.

$$\begin{array}{ll}
 x^{\frac{1}{2}} = \sqrt{x} & 2^{\frac{1}{2}} = \sqrt{2} \\
 \sqrt{x} \cdot \sqrt{x} = x & \sqrt{3} \cdot \sqrt{3} = 3 \\
 \sqrt{xy} = \sqrt{x} \cdot \sqrt{y} & \sqrt{12} = \sqrt{4 \cdot 3} = \sqrt{4} \cdot \sqrt{3} = 2 \cdot \sqrt{3} \\
 x^{\frac{1}{n}} = \sqrt[n]{x} & 5^{\frac{1}{3}} = \sqrt[3]{5} \\
 x^{\frac{m}{n}} = \sqrt[n]{x^m} & 3^{-\frac{2}{5}} = \frac{1}{\sqrt[5]{3^2}}
 \end{array}$$

1.2.3 Laws of logarithms

Logarithms are the inverse mathematical operation of exponents, like division is the inverse mathematical operation of multiplication. The logarithm is often used to find the variable in an exponent.

$$a^x = b \Leftrightarrow x = \log_a b$$

Since $\log_a a^x = x$, so then $x = \log_a b$.

This formula shows that the variable x in the power of the exponent becomes the subject of your log equation, while the number a becomes the base of your logarithm.

Logarithms with bases of 10 and e have special notations in which their base is not explicitly noted.

$$\begin{aligned}\log_{10} x &= \log x \\ \log_e x &= \ln x\end{aligned}$$

Below are the rules that you will need to use when performing calculations with logarithms and when simplifying them. The sets of equations on the left and right are the same; on the right we show the notation that the formula booklet uses while the equations on the left are easier to understand.

Laws of logarithms and change of base

I:	$\log A + \log B = \log(A \cdot B)$	$\log_a(xy) = \log_a x + \log_a y$
II:	$\log A - \log B = \log\left(\frac{A}{B}\right)$	$\log_a\left(\frac{x}{y}\right) = \log_a x - \log_a y$
III:	$n \log A = \log(A^n)$	$\log_a(x^m) = m \log_a x$
IV:	$\log_B A = \frac{\log A}{\log B}$	$\log_a x = \frac{\log_b x}{\log_b a}$

Next to these rules, there are a few handy things to keep in mind when working with logarithms.

$$\begin{aligned}\log_a 0 &= x \text{ is always undefined (because } a^x \neq 0) \\ x = \log_a a &= 1, \text{ which also means that } \ln e = 1 \\ e^{\ln a} &= a\end{aligned}$$

DB 1.5

Remember that e is just the irrational number $2.71828\dots$, so the same laws apply to \ln as to other logarithms.

DB 1.7

With the 4th rule you can change the base of a log

Solve for x in the exponent using logarithms

Solve $2^x = 13$

1. Take the log on both sides $\log 2^x = \log 13$

2. Use rule III to take x outside $x \log 2 = \log 13$

3. Solve $x = \frac{\log 13}{\log 2}$

Expressing logs in terms of other logs

Example.

Given that $p = \log_a 5$ and $q = \log_a 2$ express the following in terms of p and q

a $\log_a 10$	$\log_a(5 \times 2) = \log_a 5 + \log_a 2 = p + q$
b $\log_a 8$	$\log_a(2^3) = 3 \log_a 2 = 3q$
c $\log_a 2.5$	$\log_a\left(\frac{5}{2}\right) = \log_a 5 - \log_a 2 = p - q$

1.3 Binomial expansion



Binomial expression an expression $(a + b)^n$ which is the sum of two terms raised to the power n .

e.g. $(x + 3)^2$

Binomial expansion $(a + b)^n$ expanded into a sum of terms

e.g. $x^2 + 6x + 9$

Binomial expansions get increasingly complex as the power increases:

binomial **binomial expansion**

$$(a + b)^1 = a + b$$

$$(a + b)^2 = a^2 + 2ab + b^2$$

$$(a + b)^3 = a^3 + 3a^2b + 3ab^2 + b^3$$

The general formula for each term in the expansion is ${}^n C_r a^{n-r} b^r$.

In order to find the full binomial expansion of a binomial, you have to determine the **coefficient** ${}^n C_r$ and the **powers** for each term. The powers for a and b are found as $n - r$ and r respectively, as shown by the binomial expansion formula.

Binomial expansion formula

DB 1.9


$$\begin{aligned} (a + b)^n &= a^n + {}^n C_1 a^{n-1} b + \dots + {}^n C_r a^{n-r} b^r + \dots + b^n \\ &= {}^n C_0 a^n + {}^n C_1 a^{n-1} b + {}^n C_2 a^{n-2} b^2 + \dots \end{aligned}$$

The powers decrease by 1 for a and increase by 1 for b for each subsequent term.

The sum of the powers of each term will always = n .

There are two ways to find the coefficients: with Pascal's triangle or the binomial coefficient function (nCr). You are expected to know both methods.

Pascal's triangle



Pascal's triangle is an easy way to find all the coefficients for your binomial expansion. It is particularly useful in cases where:

1. the power is not too high (because you have to write it out manually)
2. you need to find all the terms in a binomial expansion

Binomial coefficient functions

The alternative is to calculate the individual coefficients using the nCr function on your calculator, or with the formula below.

In the 1st term of the expansion $r = 0$, in the 2nd term $r = 1, \dots$

$${}^n C_r = \frac{n!}{r!(n-r)!}$$

Expanding binomial expressions

Find the expansion of $\left(x - \frac{2}{x}\right)^5$

1. Use the binomial expansion formula

$$a = x, b = -\frac{2}{x} \text{ and } n = 5$$

$$\begin{aligned} &(x)^5 + (5C1)(x)^4\left(-\frac{2}{x}\right) + \\ &(5C2)(x)^3\left(-\frac{2}{x}\right)^2 + (5C3)(x)^2\left(-\frac{2}{x}\right)^3 + \\ &(5C4)(x)\left(-\frac{2}{x}\right)^4 + (5C5)\left(-\frac{2}{x}\right)^5 \end{aligned}$$

2. Find coefficients using Pascal's triangle for low powers or nCr on calculator for high powers

Row 0:						1
Row 1:					1	1
Row 2:				1	2	1
Row 3:			1	3	3	1
Row 4:		1	4	6	4	1
Row 5:	1	5	10	10	5	1

$(5C0)=1 \quad (5C1)=5 \quad (5C2)=10 \quad (5C3)=10 \quad (5C4)=5 \quad (5C5)=1$

3. Put the terms and their coefficients together

$$\begin{aligned} &x^5 + 5x^4\left(-\frac{2}{x}\right)^1 + 10x^3\left(-\frac{2}{x}\right)^2 + \\ &10x^2\left(-\frac{2}{x}\right)^3 + 5x\left(-\frac{2}{x}\right)^4 + \left(-\frac{2}{x}\right)^5 \end{aligned}$$



4. Simplify using laws of exponents


$$x^5 - 10x^3 + 40x - \frac{80}{x} + \frac{80}{x^3} - \frac{32}{x^5}$$

Finding a specific term in a binomial expansion

Find the coefficient of x^5 in the expansion of $(2x - 5)^8$

- | | | |
|----|---|--|
| 1. | Use the binomial expansion formula | $(a + b)^n = \dots + {}^n C_r a^{n-r} b^r + \dots$ |
| 2. | Determine r | Since $a = 2x$, to find x^5 we need a^5 .
$a^5 = a^{n-r} = a^{8-r}$, so $r = 3$ |
| 3. | Plug r into the general formula | ${}^n C_r a^{n-r} b^r = {}^8 C_3 a^{8-3} b^3 = {}^8 C_3 a^5 b^3$ |
| 4. | Replace a and b | ${}^8 C_3 (2x)^5 (-5)^3$ |
| 5. | Use nCr to calculate the value of the coefficient, ${}^n C_r$ | ${}^8 C_3 = 8C3 = 56$ |



Press 
5: Probability
3: Combinations

Insert the values for n and r separated by a comma

- | | | |
|----|-------------------------|---|
| 6. | Substitute and simplify | $56 \times 2^5 (x^5) \times (-5)^3 = -224000(x^5)$
\Rightarrow coefficient of x^5 is -224000 |
|----|-------------------------|---|

The IB use three different terms for these types of question which will effect the answer you should give.



- Coefficient** the number before the x value
- Term** the number and the x value
- Constant term** the number for which there is no x value (x^0)

FUNCTIONS

Table of contents & cheatsheet

Definitions

Function a mathematical relationship where each input has a single output. It is often written as $f(x)$ where x is the input

Domain all possible x values, the input. (the domain of investigation)

Range possible y values, the output. (the range of outcomes)

Coordinates uniquely determines the position of a point, given by (x, y)

2.1. Types of functions


22

Linear functions $y = mx + c$
 m is the *gradient*,
 c is the *y intercept*.

Midpoint: $\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}\right)$

Distance: $\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$

Gradient: $m = \frac{y_2 - y_1}{x_2 - x_1}$




Parallel lines: $m_1 = m_2$ (same gradients)

Perpendicular lines: $m_1 m_2 = -1$

Quadratic functions $y = ax^2 + bx + c = 0$

Axis of symmetry: x -coordinate of the vertex: $x = \frac{-b}{2a}$

Factorized form: $y = (x + p)(x + q)$



If $a = 1$ use the factorization method $(x + p) \cdot (x + q)$

If $a \neq 1$ use the quadratic formula


When asked explicitly complete the square

Vertex form: $y = a(x - h)^2 + k$

Vertex: (h, k)


Exponential

$$f(x) = a^x + c$$



Logarithmic


$$g(x) = \log_a(x + b)$$



2.2. Rearranging functions

28

Inverse function, $f^{-1}(x)$ reflection of $f(x)$ in $y = x$.



Composite function, $(f \circ g)(x)$ is the combined function f of g of x .


When $f(x)$ and $g(x)$ are given, replace x in $f(x)$ by $g(x)$.

Transforming functions

Change to $f(x)$	Effect
$f(x) + a$	Move graph a units upwards
$f(x + a)$	Move graph a units to the left
$a \cdot f(x)$	Vertical stretch by factor a
$f(a \cdot x)$	Horizontal stretch by factor $\frac{1}{a}$
$-f(x)$	Reflection in x -axis
$f(-x)$	Reflection in y -axis

2.1 Types of functions

Functions are mathematical relationships where each input has a single output. You have probably been doing functions since you began learning maths, but they may have looked like this:



Algebraically this is:
 $f(x) = x + 10$,
 here $x = 16, y = 26$.

We can use graphs to show multiple outputs of y for inputs x , and therefore visualize the relation between the two. Two common types of functions are linear functions and quadratic functions.

2.1.1 Linear functions



Linear functions $y = mx + c$ increases/decreases at a constant rate m , where m is the gradient and c is the y -intercept


Midpoint $\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right)$

Distance $\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}$

Gradient $m = \frac{y_2 - y_1}{x_2 - x_1}$

Parallel lines $m_1 = m_2$ (equal gradients)

Perpendicular lines $m_1 m_2 = -1$



Straight line equations are sometimes written in two other forms, which you should be comfortable rearranging them to and from:

$$ax + by + d = 0$$

general form

$$y - y_1 = m(x - x_1)$$

point-slope form

Example.

Determine the midpoint, length and gradient of the straight line connecting the two points $P_1(2, 8)$ and $P_2(6, 3)$


Midpoint: $\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2} \right) = \left(\frac{2 + 6}{2}, \frac{8 + 3}{2} \right) = (4, 5.5)$

Distance: $\sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2} = \sqrt{(6 - 2)^2 + (3 - 8)^2} = \sqrt{(4)^2 + (5)^2} = \sqrt{41}$

Gradient: $m = \frac{y_2 - y_1}{x_2 - x_1} = m = \frac{3 - 8}{6 - 2} = -\frac{5}{4}$


Parallel line: $-\frac{5}{4}x + 3$

Perpendicular line: $\frac{4}{5}x + 7$




2.1.2 Quadratic functions

Graphed, quadratic functions make a parabolic shape; they increase/decrease at an increasing rate.



$a > 0$, positive quadratic



$a < 0$, negative quadratic



Quadratic functions $y = ax^2 + bx + c = 0$

Axis of symmetry $x = \frac{-b}{2a} = x\text{-coordinate of vertex}$

Roots the x -intercept(s). Algebraically, roots can be found through factorisation or using the quadratic formula

Quadratic formula used to find the roots if $a \neq 1$

$$\frac{-b \pm \sqrt{b^2 - 4ac}}{2a} = \frac{-b \pm \sqrt{\Delta}}{2a}$$

Discriminant Δ the $b^2 - 4ac$ part of the formula, which can be used to determine how many x -intercepts a quadratic equation has

$$\Delta > 0 \Rightarrow 2 \text{ roots}$$

$$\Delta = 0 \Rightarrow 1 \text{ root}$$

$$\Delta < 0 \Rightarrow \text{no real roots}$$

DB 2.6

Roots of quadratic equations are also referred to as 'solutions' or 'zeros'.

Two more standard ways to write quadratic equations are:

$$y = a(x - h)^2 + k$$

with the vertex at (h, k)

$$y = a(x - p)(x - q)$$

with x -intercepts $(p, 0)(q, 0)$

Solving quadratic equations by factorisation

Solve $x^2 - 5x + 6 = 0$

- | | | |
|----|---|--|
| 1. | Set up a system of equations
$p + q = b$ and $p \times q = c$ | $\left. \begin{array}{l} p + q = -5 \\ p \times q = 6 \end{array} \right\} p = -2 \text{ and } q = -3$ |
| 2. | Plug the values for p and q into
$(x + p)(x + q)$ | $(x - 2)(x - 3) = x^2 - 5x + 6$ |
| 3. | Equate each part to 0
$(x + p) = 0$, $(x + q) = 0$,
and solve for x | $\left. \begin{array}{l} (x - 2) = 0 \\ (x - 3) = 0 \end{array} \right\} x = 2 \text{ or } x = 3$ |

Solving quadratic equations using the quadratic formula

Solve $3x^2 - 8x + 4 = 0$

- | | | |
|----|---|--|
| 1. | Calculate the discriminant
$\Delta = b^2 - 4ac$ | $\Delta = (-8)^2 - 4 \cdot 3 \cdot 4 = 16$ |
| 2. | Use the discriminant to determine the
number of solutions | $\Delta > 0 \text{ so 2 solutions}$ |
| 3. | Find solutions using quadratic formula
$x = \frac{-b \pm \sqrt{\Delta}}{2a}$ | $\left. \begin{array}{l} x = \frac{8 \pm \sqrt{16}}{2 \cdot 3} = \frac{8 \pm 4}{6} \\ = \frac{8 - 4}{6} = \frac{4}{6} \\ = \frac{8 + 4}{6} = 2 \end{array} \right\} \Rightarrow x = \frac{2}{3} \text{ or } x = 2$ |

Through a method called completing the square, you can rearrange a quadratic function into the form $y = a(x - b)^2 + k$. This way you can find the coordinates of the vertex (the minimum or maximum). For the exam you will always be asked explicitly to use this method.

Find the vertex by completing the square	
Express $f(x) = 4x^2 - 2x - 5 = 0$ in the form $y = a(x - b)^2 + k$. Hence, find the coordinate of the vertex of $f(x)$.	
1. Move c to the other side	$4x^2 - 2x = 5$
2. Divide by a	$x^2 - \frac{1}{2}x = \frac{5}{4}$
3. Calculate $\left(\frac{x \text{ coefficient}}{2}\right)^2$	$\left(\frac{-\frac{1}{2}}{2}\right)^2 = \frac{1}{16}$
4. Add this term to both sides	$x^2 - \frac{1}{2}x + \frac{1}{16} = \frac{5}{4} + \frac{1}{16}$
5. Factor perfect square and bring constant back	$\left(x - \frac{1}{4}\right)^2 - \frac{21}{16} = 0$ \Rightarrow minimum point = $\left(\frac{1}{4}, -\frac{21}{16}\right)$

2.1.3 Functions with asymptotes



Asymptote a straight line that a curve approaches, but never touches.

A single function can have multiple asymptotes: horizontal, vertical and in rare cases diagonal. Functions that contain the variable (x) in the denominator of a fraction and exponential and logarithmic functions will always have asymptotes.

Vertical asymptotes

Vertical asymptotes occur when the denominator is zero, as dividing by zero is undefinable. Therefore if the denominator contains x and there is a value for x for which the denominator will be 0, we get a vertical asymptote.

Example.

In the function $f(x) = \frac{x}{x-4}$ the denominator is 0 when $x = 4$, so this line forms the a vertical asymptote.

Horizontal asymptotes

Horizontal asymptotes are the value that a function tends to as x becomes really big or really small; technically speaking to the limit of infinity, $x \rightarrow \infty$. The general idea is then that when x is large, other parts of the function not involving x become insignificant and so can be ignored.

Example.

In the function $f(x) = \frac{x}{x-4}$, when x is small the 4 is important.

$$x = 10 \qquad 10 - 4 = 6$$

But as x gets bigger the 4 becomes increasingly insignificant

$$x = 100 \qquad 100 - 4 = 96$$

$$x = 10000 \qquad 10000 - 4 = 9996$$

Therefore as we approach the limits we can ignore the 4.

$$\lim_{x \rightarrow \infty} f(x) = \frac{x}{x} = 1$$


So there is a horizontal asymptote at $y = 1$.

Exponential and logarithmic functions

Exponential functions will always have a horizontal asymptote and logarithmic functions will always have a vertical asymptote, due to the nature of these functions. The position of the asymptote is determined by constants in the function.


Exponential

$f(x) = a^x + c$ asymptote at $y = c$
 where a is a positive number (often e)



Logarithmic

$g(x) = \log_a(x + b)$ asymptote at $x = -b$




2.1.4 Special Functions

The function $\frac{ax + b}{cx + d}$



Rational function of the form $y = \frac{ax + b}{cx + d}$

The graph made by this function has one horizontal and one vertical asymptote. The graph is not continuous in all points, but splits into two parts. Both of the parts approach horizontal asymptote at either negative or positive large values of x and approach horizontal asymptote at either negative or positive large values of y . Also both parts are located in different “corners” of the coordinate system. The vertical asymptote occurs where denominator is equal to 0, meaning $x = -\frac{d}{c}$. The graph is not defined at that point.




The horizontal asymptote occurs at very large values of x , meaning that asymptote occurs at

$$y = \lim_{x \rightarrow \infty} \frac{ax + b}{cx + d} = \frac{a}{c}$$


2.2 Rearranging functions

2.2.1 Inverse functions, $f^{-1}(x)$

Inverse functions are the reverse of a function. Finding the input x for the output y . You can think of it as going backwards through the number machine.



This is the same as reflecting a graph in the $y = x$ axis.



Finding the inverse function.

$$f(x) = 2x^3 + 3, \text{ find } f^{-1}(x)$$

- | | | |
|----|---|--|
| 1. | Replace $f(x)$ with y | $y = 2x^3 + 3$ |
| 2. | Solve for x | $y - 3 = 2x^3$ $\Rightarrow \frac{y - 3}{2} = x^3$ $\Rightarrow \sqrt[3]{\frac{y - 3}{2}} = x$ |
| 3. | Replace x with $f^{-1}(x)$ and y with x | $\sqrt[3]{\frac{x - 3}{2}} = f^{-1}(x)$ |

2.2.2 Composite functions

Composite functions are a combination of two functions.

$$(f \circ g)(x) \quad \text{means } f \text{ of } g \text{ of } x$$

To find the composite function above substitute the function of $g(x)$ into the x of $f(x)$.

Let $f(x) = 2x + 3$ and $g(x) = x^2$. Find $(f \circ g)(x)$ and $(g \circ f)(x)$.

$(f \circ g)(x)$: replace x in the $f(x)$ function with the entire $g(x)$ function

$$(2g(x)) + 3 = 2x^2 + 3$$

$(g \circ f)(x)$: replace x in the $g(x)$ function with the entire $f(x)$ function

$$(f(x))^2 = (2x + 3)^2$$

Remember
 $f \circ g(x) \neq g \circ f(x)$

Example.

2.2.3 Transforming functions

By adding and/or multiplying by constants we can transform a function into another function.

Change to $f(x)$	Effect
$a \cdot f(x)$	Vertical stretch by factor a
$f(a \cdot x)$	Horizontal stretch by factor $1/a$
$-f(x)$	Reflection in x -axis
$f(-x)$	Reflection in y -axis
$f(x) + a$	Move graph a units upwards
$f(x + a)$	Move graph a units to the left


Exam hint: describe the transformation with words as well to guarantee marks.

Always do translations last


Transforming functions $f(x) \rightarrow af(x + b)$

Given $f(x) = \frac{1}{4}x^3 + x^2 - \frac{5}{4}x$, draw $3f(x - 1)$.

1. Sketch $f(x)$




2. Stretch the graph by the factor of a



3. Move graph by $-b$

Move graph by 1 to the right






2.3 Intersection

When functions intersect the x and y -values are equal, so at the point of intersection $f(x) = g(x)$.

To find the intersection point(s) of two functions.

$f(x) = \left(\frac{1}{2}\right)^x - 2$ and $g(x) = -x^2 + 4$. Find the coordinates where $f(x) = g(x)$.



Plot both functions

Press on , go to "Graph"

Enter the two functions:

$$f1(x) = \left(\frac{1}{2}\right)^x - 2,$$

press tab to input

$$f2(x) = -x^2 + 4$$

Find the intersection:

Press menu

8: Geometry

1: Points & lines

3: Intersection Point(s)

Approach the intersection

you are trying to find with the cursor and click once you near it. Repeat for any other intersections.

In this case the intersection points are $(-1.68, 1.19)$ and $(2.41, -1.81)$.

Note: if you can't find 8: Geometry make sure you use graph mode instead of the scratchpad, otherwise please update your N-spire to the latest version.

TRIGONOMETRY AND CIRCULAR FUNCTIONS


Table of contents & cheatsheet

3.2. Basic trigonometry

33

$$\text{radians} = \frac{\pi}{180^\circ} \times \text{degrees} \quad \text{degrees} = \frac{180^\circ}{\pi} \times \text{radians}$$

Before each question make sure calculator is in correct setting: degrees or radians?




$$\text{Area of a sector} = \frac{1}{2} r^2 \cdot \theta$$

$$\text{Arc length} = r \cdot \theta$$

θ in radians, r = radius.

Right-angle triangle (triangle with 90° angle)



$$\sin \theta = \frac{\text{opposite}}{\text{hypotenuse}} \quad \text{SOH}$$


$$\cos \theta = \frac{\text{adjacent}}{\text{hypotenuse}} \quad \text{CAH}$$

$$\tan \theta = \frac{\text{opposite}}{\text{adjacent}} \quad \text{TOA}$$

Three-figure bearings

Direction given as an angle of a full circle. North is 000° and the angle is expressed in the clockwise direction from North. So East is 090° , South is 180° and West 270° .

Non-right angle triangles



$$\text{Sine rule: } \frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$$

Use this rule when you know: 2 angles and a side (not between the angles) or 2 sides and an angle (not between the sides).

$$\text{Cosine rule: } c^2 = a^2 + b^2 - 2ab \cos C$$


Use this rule when you know: 3 sides or 2 sides and the angle between them.

$$\text{Area of a triangle: } \text{Area} = \frac{1}{2} ab \sin C$$

Use this rule when you know: 3 sides or 2 sides and the angle between them.

3.3. Circular functions

39



deg	0°	30°	45°	60°	90°	120°	135°	150°	180°
rad	0	$\frac{1}{6}\pi$	$\frac{1}{4}\pi$	$\frac{1}{3}\pi$	$\frac{1}{2}\pi$	$\frac{2}{3}\pi$	$\frac{3}{4}\pi$	$\frac{5}{6}\pi$	π
$\sin \theta$	0	$\frac{1}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{\sqrt{3}}{2}$	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0
$\cos \theta$	1	$\frac{\sqrt{3}}{2}$	$\frac{\sqrt{2}}{2}$	$\frac{1}{2}$	0	$-\frac{1}{2}$	$-\frac{\sqrt{2}}{2}$	$-\frac{\sqrt{3}}{2}$	-1
$\tan \theta$	0	$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	∞	$-\sqrt{3}$	-1	$-\frac{1}{\sqrt{3}}$	0

Trigonometric function $y = a \sin(bx + c) + d$

Amplitude: a

Period: $\frac{360^\circ}{b}$ or $\frac{2\pi}{b}$

Horizontal shift: c

Vertical shift: d

Trigonometric identities

$$\tan \theta = \frac{\sin \theta}{\cos \theta}$$

$$\sin^2 \theta + \cos^2 \theta = 1$$

$$2 \sin \theta \cos \theta = \sin 2\theta$$

$$\cos 2\theta = \cos^2 \theta - \sin^2 \theta$$

3.1 Properties of 3D shapes

3.1.1 Points in 3D space

When you are learning about the points on a 2-dimensional plane, you also learn how to find distances between those two points. One of the easiest ways to derive that formula is by constructing a triangle and using Pythagoras. In the same way it is possible to derive a very similar expression for distance between two points in a 3D space:

$$d = \sqrt{(x_1 - x_2)^2 + (y_1 - y_2)^2 + (z_1 - z_2)^2}$$





You have also learnt how to find the midpoint between the two points (x_1, y_1) and (x_2, y_2) : add those individual coordinates together and divide the sum by two. One can find the midpoint between two points in 3D space in almost exact same way:

$$\left(\frac{x_1 + x_2}{2}, \frac{y_1 + y_2}{2}, \frac{z_1 + z_2}{2} \right)$$

3.1.2 Pyramids, cones and spheres

In the exam you might be asked to find volume or surface of different 3D objects. These objects might be either familiar to you 3D shapes or made up from those shapes. In the first case, the formulas should be given in the data booklet. In the latter case, you would need to split the object into familiar shapes and sum the required components together.

DB 3.1

Volume of a right-pyramid	$V = \frac{1}{3}Ab$	
Volume of a right cone	$V = \frac{1}{3}\pi r^2 h$	
Area of the curved surface of a cone	$A = \pi r l$	
Volume of a sphere	$V = \frac{4}{3}\pi r^3$	
Surface area of a sphere	$A = 4\pi r^2$	

3.2 Basic trigonometry

This section offers an overview of some basic trigonometry rules and values that will recur often. It is worthwhile to know these by heart; but it is much better to understand how to obtain these values. Like converting between Celsius and Fahrenheit; you can remember some values that correspond to each other but if you understand how to obtain them, you will be able to convert any temperature.

3.2.1 Converting between radians and degrees

$$\text{radians} = \frac{\pi}{180^\circ} \times \text{degrees}$$

$$\text{degrees} = \frac{180^\circ}{\pi} \times \text{radians}$$




Table 3.1: Common radians/degrees conversions


Degrees	0°	30°	45°	60°	90°	120°	135°	180°	270°	360°
Radians	0	$\frac{\pi}{6}$	$\frac{\pi}{4}$	$\frac{\pi}{3}$	$\frac{\pi}{2}$	$\frac{2\pi}{3}$	$\frac{3\pi}{4}$	π	$\frac{3\pi}{2}$	2π

3.2.2 Circle formulas

$$\text{Area of a sector} = \frac{1}{2}r^2 \cdot \vartheta$$

$$\text{Arc length} = r \cdot \vartheta$$


ϑ in radians, r = radius.





DB 3.4

3.2.3 Right-angle triangles



$a^2 = b^2 + c^2$	Pythagoras
$\sin \vartheta = \frac{\text{opposite}}{\text{hypotenuse}}$	SOH
$\cos \vartheta = \frac{\text{adjacent}}{\text{hypotenuse}}$	CAH
$\tan \vartheta = \frac{\text{opposite}}{\text{adjacent}}$	TOA




The following two right angle triangles with whole numbers for all the sides come up often in past exam questions.



The two triangles below can help you in finding the sin, cos and tan of the angles that you should memorize, shown in table 3.2 on page 40. Use SOH, CAH, TOA to find the values.



3.2.4 Non-right angle triangles



To find any missing angles or side lengths in non-right angle triangles, use the *cosine* and *sine* rule. Remember that the angles in the triangle add up to 180°!


Read the question: does it specify if you are looking for an acute (less than 90°) or obtuse (more than 90°) angle? If not there may be 2 solutions. Exam hint: Use sketches when working with worded questions!

DB 3.2

Sine rule: $\frac{a}{\sin A} = \frac{b}{\sin B} = \frac{c}{\sin C}$


Use this rule when you know:

2 angles and a side
(not between the angles)



or


2 sides and an angle
(not between the sides)



Cosine rule: $c^2 = a^2 + b^2 - 2ab \cos C$


Use this rule when you know:

3 sides



or


2 sides and the angle between them



Area of a triangle: $\text{Area} = \frac{1}{2}ab \sin C$

Use this rule when you know:

2 sides and the angle between them



Example.

$\triangle ABC: A = 40^\circ, B = 73^\circ, a = 27 \text{ cm}.$

Find $\angle C$

$$\angle C = 180^\circ - 40^\circ - 73^\circ = 67^\circ$$

Find b

$$\begin{aligned} \frac{a}{\sin A} &= \frac{b}{\sin B} \\ \frac{27}{\sin 40^\circ} &= \frac{b}{\sin 73^\circ} \\ b &= \frac{27}{\sin 40^\circ} \cdot \sin 73^\circ = 40.169 \approx 40.2 \text{ cm} \end{aligned}$$


Find c

$$\begin{aligned} \frac{c}{\sin C} &= \frac{a}{\sin A} \\ c &= \frac{27}{\sin 40^\circ} \times \sin 67^\circ = 38.7 \text{ cm} \end{aligned}$$

Find the area

$$\begin{aligned} \text{Area} &= \frac{1}{2} \cdot 27 \cdot 40 \cdot \sin 67^\circ \\ &= 499.59 \approx 500 \text{ cm}^2 \end{aligned}$$

Example.



Find z

$$\begin{aligned} z^2 &= 6^2 + 10^2 - 2 \cdot 6 \cdot 10 \cdot \cos 35^\circ \\ z^2 &= 37.70 \\ z &= 6.14 \text{ km} \end{aligned}$$

Find $\angle x$

$$\begin{aligned} \frac{6}{\sin x} &= \frac{6.14}{\sin 35^\circ} \\ \sin x &= 0.56 \\ x &= \sin^{-1}(0.56) = 55.91^\circ \end{aligned}$$

3.2.5 Ambiguous case

Ambiguous case, also known as an angle-side-side case, is when the triangle is not unique from the given information. It happens when you are given two sides and an angle not between those sides in a triangle.

You have to use a sine rule to solve a problem in this case. However, one needs to remember that $\sin x = \sin(180^\circ - x)$, meaning that your answer for an angle is not just x , but also $180^\circ - x$.

In other words, we might get two different possible angles as an answer and thus two different possible triangles that satisfy the information given.

However, that is not always the case, if the sum of the two known angles becomes bigger than 180° . So if you are required to calculate the third angle or total area of a triangle, you might have to do the calculations for two different triangles using both of your angles.



Example.

$\triangle ABC : B = 33^\circ, a = 23 \text{ cm}, b = 14 \text{ cm}.$


Find $\angle A$.

$$\begin{aligned} \frac{a}{\sin A} &= \frac{b}{\sin B} \\ \frac{23}{\sin 33^\circ} &= \frac{14}{\sin A} \\ \angle A_1 &= 63.5^\circ \\ \angle A_2 &= 180^\circ - 63.5^\circ = 117^\circ \\ \angle A_2 + 33^\circ &< 180^\circ \text{ thus also a possible angle} \end{aligned}$$

Draw the two possible triangles.



3.2.6 Three-figure bearings




Three-figure bearings can be used to indicate compass directions on maps. They will be given as an angle of a full circle, so between 0° and 360° . North is always marked as 0° . Any direction from there can be expressed as the angle in the clockwise direction from North.


In questions on three-figure bearings, you are often confronted with quite a lot of text, so it is a good idea to first make a drawing. You may also need to create a right angle triangle and use your basic trigonometry.

Example.

SW: 45° between South and West = 225°




N 40° E: 40° East of North = 040°



Solving questions with three-figure bearings

A ship left port A and sailed 20 km in the direction 120° . It then sailed north for 30 km to reach point C . How far from the port is the ship?

1. Sketch



2. Find an internal angle of the triangle

$$\vartheta = 180^\circ - 120^\circ = 60^\circ = C$$

Similar angles between two parallel lines

3. Use cosine or sin rule

(here cosine)

$$AC^2 = AB^2 + BC^2 - 2 \cdot AB \cdot BC \cdot \cos \vartheta$$


$$AC^2 = 20^2 + 30^2 - 2 \cdot 20 \cdot 30 \cdot \cos 60^\circ$$

$$AC^2 = 400 + 900 - 2 \cdot 20 \cdot 30 \cdot \frac{1}{2}$$

$$AC = \sqrt{400 + 900 - 600} = \sqrt{700}$$

3.3 Circular functions


3.3.1 Unit circle



The unit circle is a circle with a radius of 1 drawn from the origin of a set of axes. The y -axis corresponds to *sine* and the x -axis to *cosine*; so at the coordinate $(0, 1)$ it can be said that $\cosine = 0$ and $sine = 1$, just like in the $\sin x$ and $\cos x$ graphs when plotted.

The unit circle is a tool that you can use when solving problems involving circular functions. You can use it to find all the solutions to a trigonometric equation within a certain domain.

As you can see from their graphs, functions with $\sin x$, $\cos x$ or $\tan x$ repeat themselves every given period; this is why they are also called *circular functions*. As a result, for each y -value there is an infinite amount of x -values that could give you the same output. This is why questions will give you a set domain that limits the x -values you should consider in your calculations or represent on your sketch (e.g. $0^\circ \leq x \leq 360^\circ$).




Relations between \sin , \cos and \tan :


- α and β have the same sine
- α and ϑ have the same cosine
- β and ϑ have the same tangent

Example.

$\sin 30^\circ = \sin 150^\circ$



$\cos 30^\circ = \cos 330^\circ$



$\tan 150^\circ = \tan 330^\circ$





Table 3.2: Angles to memorize

deg	0°	30°	45°	60°	90°	120°	135°	150°	180°
rad	0	$\frac{1}{6}\pi$	$\frac{1}{4}\pi$	$\frac{1}{3}\pi$	$\frac{1}{2}\pi$	$\frac{2}{3}\pi$	$\frac{3}{4}\pi$	$\frac{5}{6}\pi$	π
$\sin \vartheta$	0	$\frac{1}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{\sqrt{3}}{2}$	1	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2}$	0
$\cos \vartheta$	1	$\frac{\sqrt{3}}{2}$	$\frac{1}{\sqrt{2}}$	$\frac{1}{2}$	0	$-\frac{1}{2}$	$-\frac{1}{\sqrt{2}}$	$-\frac{\sqrt{3}}{2}$	-1
$\tan \vartheta$	0	$\frac{1}{\sqrt{3}}$	1	$\sqrt{3}$	∞	$-\sqrt{3}$	-1	$-\frac{1}{\sqrt{3}}$	0


3.3.2 Graphs of trigonometric functions

sin x




Domain: $x \in \mathbb{R}$
 Amplitude: $-1 \leq y \leq 1$
 Period: $2n\pi, n \cdot 360^\circ$, with $n \in \mathbb{Z}$

cos x



Domain: $x \in \mathbb{R}$
 Amplitude: $-1 \leq y \leq 1$
 Period: $2n\pi, n \cdot 360^\circ$, with $n \in \mathbb{Z}$

tan x



Domain: $x \in \mathbb{R}, x \neq \frac{\pi}{2} + k\pi$, with $k \in \mathbb{Z}$
 Amplitude: $-\infty < y < \infty$
 Period: $n\pi, n \cdot 180^\circ$, with $n \in \mathbb{Z}$

3.3.3 Transformations

Besides the transformations in the functions chapter, trigonometric functions have some transformations with their own particular names. For a trigonometric function, the vertical stretch on a graph is determined by its amplitude, the horizontal stretch by its period and an upward/downward shift by its axis of oscillation.





A trigonometric function, given by $y = a \sin(bx + c) + d$, has:

- amplitude a
- period of $\frac{360^\circ}{b}$ or $\frac{2\pi}{b}$
- horizontal shift of $+c$ to the left, in degrees or radians
- vertical shift of $+d$ upwards, oscillates around d .

A negative a will flip your graph around the x -axis. Negative values of c and d will lead to shifts to the right and downwards the respective number of units

Example

Transformations of $y = \cos x$.



3.3.4 Identities and equations

DB 3.5 & 3.6

In order to solve trigonometric equations, you will sometimes need to use identities. Identities allow you to rewrite your equation in a way that will make it easier to solve algebraically.

Trigonometric identity

$$\tan \vartheta = \frac{\sin \vartheta}{\cos \vartheta}$$

$$\sin^2 \vartheta + \cos^2 \vartheta = 1$$

$$2 \sin \vartheta \cos \vartheta = \sin 2\vartheta$$

$$\cos 2\vartheta = \cos^2 \vartheta - \sin^2 \vartheta$$

Popular rearrangement

$$\cos \vartheta = \frac{\sin \vartheta}{\tan \vartheta}$$

$$\sin \vartheta = \cos \vartheta \times \tan \vartheta$$

$$\sin^2 \vartheta = 1 - \cos^2 \vartheta$$

$$\cos^2 \vartheta = 1 - \sin^2 \vartheta$$

$$\cos 2\vartheta = 1 - 2 \sin^2 \vartheta$$

$$\cos 2\vartheta = 2 \cos^2 \vartheta - 1$$

Solving equations with trigonometric identities

Solve $2 \cos^2 x + \sin x = 1$, $0^\circ \leq x \leq 360^\circ$.

- Identify which identity from the formula booklet to use. Note that you are always aiming to get an equation with just, sin, cos or tan.

Here we could use either $\sin^2 \vartheta + \cos^2 \vartheta = 1$ or $\cos^2 \vartheta - \sin^2 \vartheta = \cos 2\vartheta$. We will use the first so that we get an equation with just sin.

- Rearrange identity and substitute into equation.

$$\cos^2 \vartheta = 1 - \sin^2 \vartheta$$

$$2(1 - \sin^2 x) + \sin x = 1$$

$$2 - 2 \sin^2 x + \sin x = 1$$

$$-2 \sin^2 x + \sin x + 1 = 0$$

- Solve for x giving answers within the stated range. Recognise that here the equation looks like a quadratic equation.

Substitute u for $\sin x$:

$$-2u^2 + u + 1 = 0$$

$$(-2u - 1)(u - 1) = 0$$

$$u = \sin x \Rightarrow 1 \quad x \Rightarrow 90^\circ$$

$$u = \sin x \Rightarrow -0.5 \quad x \Rightarrow 210^\circ \text{ or } 330^\circ$$

DIFFERENTIATION

Table of contents & cheatsheet

Definitions

Differentiation is a way to find the gradient of a function at any point, written as $f'(x)$, y' and $\frac{dy}{dx}$.

Tangent line to a point on a curve is a linear line with the same gradient as that point on the curve.

4.2. Tangent and normal

46

Tangent line with the same gradient as a point on a curve.

Normal perpendicular to the tangent $m = \frac{-1}{\text{slope of tangent}}$

Both are linear lines with general formula: $y = mx + c$.

1. Use derivative to find gradient of the tangent. For normal then do $-\frac{1}{\text{slope of tangent}}$.
2. Input the x -value of the point into $f(x)$ to find y .
3. Input y , m and the x -value into $y = mx + c$ to find c .

4.4. Sketching graphs

50

Gather information before sketching:

Intercepts x -intercept: $f(x) = 0$
 y -intercept: $f(0)$

Turning points minima: $f'(x) = 0$ and $f''(x) < 0$
 maxima: $f'(x) = 0$ and $f''(x) > 0$
 point of inflection: $f''(x) = 0$

Asymptotes vertical: x -value when the function divides by 0
 horizontal: y -value when $x \rightarrow \infty$

Plug the found x -values into $f(x)$ to determine the y -values.

4.1. Polynomials

44


Product $y = uv$, then: $y' = uv' + u'v$

Quotient $y = \frac{u}{v}$, then: $y' = \frac{vu' - uv'}{v^2}$

Chain $y = g(u)$ where $u = f(x)$, then:
 $\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$

4.3. Turning points

47




	$f'(x)$	$f''(x)$
Local minimum	0	+
Local maximum	0	-
Points of inflection		0

4.5. Applications

52

Kinematics

Derivative represents the rate of change, integration the reverse.



4.1 Polynomials

As you have learnt in the section on functions, a straight line graph has a gradient. This gradient describes the rate at which the graph is changing and thanks to it we can tell how steep the line will be. In fact gradients can be found for any function - the special thing about linear functions is that their gradient is always the same (given by m in $y = mx + c$). For polynomial functions the gradient is always changing. This is where calculus comes in handy; we can use differentiation to derive a function using which we can find the gradient for any value of x .

Using the following steps, you can find the derivative function ($f'(x)$) for any polynomial function ($f(x)$).



Polynomial a mathematical expression or function that contains several terms often raised to different powers

$$\text{e.g. } y = 3x^2, \quad y = 121x^5 + 7x^3 + x \quad \text{or} \quad y = 4x^{\frac{2}{3}} + 2x^{\frac{1}{3}}$$

Principles $y = f(x) = ax^n \quad \Rightarrow \quad \frac{dy}{dx} = f'(x) = nax^{n-1}$.

The (original) function is described by y or $f(x)$, the derivative (gradient) function is described by $\frac{dy}{dx}$ or $f'(x)$.

Derivative of a constant (number) 0

$$\text{e.g. For } f(x) = 5, \quad f'(x) = 0$$

Derivative of a sum sum of derivatives.

If a function you are looking to differentiate is made up of several summed parts, find the derivatives for each part separately and then add them together again.

$$\text{e.g. } f(x) = ax^n \quad \text{and} \quad g(x) = bx^m$$

$$f'(x) + g'(x) = nax^{n-1} + mbx^{m-1}$$

4.1.1 Rules

With more complicated functions, in which several functions are being multiplied or divided by one another (rather than just added or subtracted), you will need to use the product or quotient rules.

DB 5.6

Product rule

When functions are *multiplied*: $y = uv$

then: $y' = uv' + u'v$

which is the same as $\frac{dy}{dx} = u \frac{dv}{dx} + v \frac{du}{dx}$

Quotient rule

When functions are *divided*: $y = \frac{u}{v}$

then: $y' = \frac{vu' - uv'}{v^2}$

which is the same as $\frac{dy}{dx} = \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2}$

Example.

Let $y = x^2 \cos x$, then

$$y' = x^2(\cos x)' + (x^2)' \cos x$$

$$= -x^2 \sin x + 2x \cos x$$

Let $y = \frac{x^2}{\cos x}$, then

$$y' = \frac{(x^2)' \cos x - x^2(\cos x)'}{(\cos x)^2}$$

$$= \frac{2x \cos x + x^2 \sin x}{\cos^2 x}$$

Chain rule

When a function is inside another function: $y = g(u)$ where $u = f(x)$

then: $\frac{dy}{dx} = \frac{dy}{du} \cdot \frac{du}{dx}$.

A function inside another function is a composite function, $f \circ g(x)$, which we discussed in the Functions chapter

Differentiating with the chain rule

Let $y = (\cos x)^2$, determine the derivative y'

1. Determine what the inside (u) and outside (y) functions are

Inside function: $u = \cos x$
Outside function: $y = u^2$

2. Find u' and y'

$$u' = \frac{du}{dx} = -\sin x; \quad y' = \frac{dy}{du} = 2u$$

3. Fill in chain rule formula

$$\begin{aligned} \frac{dy}{dx} &= \frac{dy}{du} \cdot \frac{du}{dx} \\ &= 2u(-\sin x) \\ &= -2 \sin x \cos x \end{aligned}$$

4.2 Tangent and normal equation



Tangent a straight line that touches a curve at one single point. At that point, the gradient of the curve is equal to the gradient of the tangent.

Normal a straight line that is perpendicular to the tangent line:

$$\text{slope of normal} = \frac{-1}{\text{slope of tangent}}$$

For any questions with tangent and/or normal lines, use the steps described in the following example.

Finding the linear function of the tangent.

Let $f(x) = x^3$. Find the equation of the tangent at $x = 2$

- | | | |
|----|--|---|
| 1. | Find the derivative and fill in value of x to determine slope of tangent | $f'(x) = 3x^2$
$f'(2) = 3 \cdot 2^2 = 12$ |
| 2. | Determine the y value | $f(x) = 2^3 = 8$ |
| 3. | Plug the slope m and the y value in $y = mx + c$ | $8 = 12x + c$ |
| 4. | Fill in the value for x to find c | $8 = 12(2) + c \Rightarrow c = -16$
eq. of tangent: $y = 12x - 16$ |

Steps 1, 2 and 4 are identical for the equation of the tangent and normal

Finding the linear function of the normal.


Let $f(x) = x^3$. Find the equation of the normal at $x = 2$


- | | | |
|----|---|---|
| 1. | _____ | $f'(2) = 12$ |
| 2. | _____ | $f(x) = 8$ |
| 3. | Determine the slope of the normal
$m = \frac{-1}{\text{slope tangent}}$ and plug it and the y -value into $y = mx + c$ | $m = \frac{-1}{12}$
$8 = -\frac{1}{12}x + c$ |
| 4. | Fill in the value for x to find c | $8 = -\frac{1}{12}(2) + c \Rightarrow c = \frac{49}{6}$
eq. of normal: $y = -\frac{1}{12}x + \frac{49}{6}$ |


Steps 1, 2 and 4 are identical for the equation of the tangent and normal


To find the gradient of a function for any value of x .


$f(x) = 5x^3 - 2x^2 + x$. Find the gradient of $f(x)$ at $x = 3$.




Press 
 4: Calculus
 1: Numerical Derivative
 at a Point



Enter the variable used in
 your function (x) and the
 value of x that you want to
 find. Keep the settings on
 1st Derivative
 Press 



Type in your function
 press 

In this case, $f'(3) = 124$

4.3 Turning points

There are three types of turning points:


1. **Local maxima**
2. **Local minima**
3. **Points of inflection**

We know that when $f'(x) = 0$ there will be a maximum or a minimum. Whether it is a maximum or minimum should be evident from looking at the graph of the original function. If a graph is not available, we can find out by plugging in a slightly smaller and slightly larger value than the point in question into $f'(x)$. If the smaller value is negative and the larger value positive then it is a local minimum. If the smaller value is positive and the larger value negative then it is a local maximum.


If you take the derivative of a derivative function (one you have already derived) you get the *second derivative*. In mathematical notation, the second derivative is written as y'' , $f''(x)$ or $\frac{d^2y}{dx^2}$. We can use this to determine whether a point on a graph is a maximum, a minimum or a point of inflection as demonstrated in the following Figure 4.1.

Figure 4.1: Graph that shows a local maximum, a local minimum and points of inflection

Notice how the points of inflection of $f(x)$ are minima and maxima in $f'(x)$ and thus equal 0 in $f''(x)$



Finding turning points



The function $f(x) = x^3 + x^2 - 5x - 5$ is shown. Use the first and second derivative to find its turning points: the minima, maxima and points of inflection (POI).

1. Find the first and second derivative

$$f'(x) = 3x^2 + 2x - 5$$

$$f''(x) = 6x + 2$$
2. Find x_{\min} and x_{\max} by setting $f'(x) = 0$

$$3x^2 + 2x - 5 = 0$$
 GDC yields: $x = 1$ or $x = -\frac{5}{3}$
3. Find y -coordinates by inserting the x -value(s) into the original $f(x)$

$$f(1) = (1)^3 + (1)^2 - 5(1) - 5 = -8,$$
 so x_{\min} at $(1, -8)$.

$$f\left(-\frac{5}{3}\right) = \left(-\frac{5}{3}\right)^3 + \left(-\frac{5}{3}\right)^2 - 5\left(-\frac{5}{3}\right) - 5 = 1.48(3 \text{ s.f.}),$$
 so x_{\max} at $\left(-\frac{5}{3}, 1.48\right)$.
4. Find POI by setting $f''(x) = 0$


$$6x + 2 = 0$$
5. Enter x -values into original function to find coordinates


$$f\left(-\frac{1}{3}\right) = \left(-\frac{1}{3}\right)^3 + \left(-\frac{1}{3}\right)^2 - 5\left(-\frac{1}{3}\right) - 5$$


$$y = -3.26(3 \text{ s.f.})$$
 so POI at $\left(-\frac{1}{3}, -3.26\right)$

Finding turning points (local maximum/minimum) of a function using GDC


Find the coordinates of the local minimum for $f(x) = 4x^2 - 5x + 3$



Press 
 6: Analyze graph
 or 2: Minimum
 or 3: Maximum



Use the cursor to set the bounds (the min/max must be between the bounds)



So the coordinates of the minimum for $f(x)$ are $(0.625, 1.44)$

4.4 Sketching graphs

When sketching a graph, you will need the following information:

1. Intercepts,
2. Turning points (maximums, minimums and inflection points) and
3. Asymptotes

Sketching a function

Sketch the function $f(x) = \frac{x^2}{x^2 - 16}$

1. Note down all information:

1. Intercepts:

- y-intercept: $f(0)$
- x-intercept: $f(x) = 0$

1. y-intercept when $x = 0$:

$$f(0) = \frac{0^2}{0^2 - 16} = 0 \quad (0, 0)$$

$$f(x) = \frac{x^2}{x^2 - 16} = 0 \quad x = 0 \quad (\text{same})$$

This is the only x-intercept.

2. Turning points:

- min/max: $f'(x) = 0$
- inflection: $f''(x) = 0$

2. Turning point: $f'(x) = \frac{-32x}{x^2 - 16^2}$,

$x = 0$ (0, 0) (Found with quotient rule).

$$f' = 0 \text{ when } x = 0.$$

3. Asymptotes:

- vertical: denominator = 0, $x = -b$, for $\log(x + b)$

- horizontal: $\lim_{x \rightarrow \infty \text{ or } x \rightarrow -\infty} a^x + c, y = c$, for

$$a^x + c$$

3. Vertical asymptotes when

$$x^2 - 16 = 0, \text{ so } x = 4 \text{ and } x = -4.$$


Horizontal asymptote:

$$\lim_{x \rightarrow \infty} f(x) = \frac{x^2}{x^2} = 1, \text{ so } y = 1$$


To find the y-coordinate, input the x-value into the original $f(x)$.

2. Mark out information on axis


Clearly indicate them to guarantee marks



3. Think about where your lines are coming from



4. Join the dots



4.5 Applications

4.5.1 Kinematics

Kinematics deals with the movement of bodies over time. When you are given one function to calculate displacement, velocity or acceleration you can use differentiation or integration to determine the functions for the other two.

$$\begin{array}{ccc}
 \frac{ds}{dt} & \left(\begin{array}{c} \text{Displacement, } s \\ \text{Velocity, } v = \frac{ds}{dt} \end{array} \right) & \int v \, dt \\
 \frac{dv}{dt} & \left(\begin{array}{c} \text{Acceleration,} \\ a = \frac{dv}{dt} = \frac{d^2s}{dt^2} \end{array} \right) & \int a \, dt
 \end{array}$$

The derivative represents the rate of change, i.e. the gradient of a graph. So, velocity is the rate of change in displacement and acceleration is the rate of change in velocity.

Answering kinematics questions.

A diver jumps from a platform at time $t = 0$ seconds. The distance of the diver above water level at time t is given by $s(t) = -4.9t^2 + 4.9t + 10$, where s is in metres. Find when velocity equals zero. Hence find the maximum height of the diver.

- | | | |
|----|--|---|
| 1. | Find an equation for velocity by differentiating equation for distance | $v(t) = -9.8t + 4.9$ |
| 2. | Solve for $v(t) = 0$ | $-9.8t + 4.9 = 0, \quad t = 0.5$ |
| 3. | Put value into equation for distance to find height above water | $s(0.5) = -4.9(0.5)^2 + 4.9(0.5) + 10 = 11.225 \text{ m}$ |

4.5.2 Optimization

We can use differentiation to find minimum and maximum areas/volumes of various shapes. Often the key skill with these questions is to find an expression using simple geometric formulas and rearranging in order to differentiate.

Finding the minimum/maximum area or volume

The sum of height and base of a triangle is 40 cm. Find an expression for its area in terms of x , its base length. Hence find its maximum area.

1.	Find expressions for relevant dimensions of the shape	$\text{length of the base } (b) = x$ $\text{height} + \text{base} = 40$ $\text{so } h + x = 40$ $\text{area of triangle } A = \frac{1}{2}xb$
2.	Reduce the number of variables by solving the simultaneous equations	$\text{Since } h = 40 - x, \text{ substitute } h \text{ into } A:$ $A = \frac{1}{2}x(40 - x) = -\frac{1}{2}x^2 + 20x$
3.	Differentiate	$f'(x) = -x + 20$
4.	Find x when $f'(x) = 0$	$-x + 20 = 0 \Rightarrow x = 20$
5.	Plug x value in $f(x)$	$-\frac{1}{2}20^2 + 20(20) = -200 + 400 = 200 \text{ cm}^2$

If an expression is given in the problem, skip to step 2 (e.g. cost/profit problems)

INTEGRATION

Table of contents & cheatsheet

5.1. Indefinite integral

56

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C \quad n \neq -1$$

Integration with an internal function

$$\int f(ax+b) dx$$

Integrate normally and multiply by $\frac{1}{\text{coefficient of } x}$

Integration by substitution

$$\int f(g(x)) \cdot g'(x) dx$$

5.2. Definite integral

58

$$\int_a^b f(x) dx = F(b) - F(a) \quad \text{where} \quad F = \int f(x) dx$$


Be careful, the order you substitute a and b into the indefinite integral is relevant for your answer:

$$\int_a^b f(x) dx = - \int_b^a f(x) dx$$

Area between a curve and the x -axis

By determining a definite integral for a function, you can find the area beneath the curve that is between the two x -values indicated as its limits.

$$A_{\text{curve}} = \int_a^b f(x) dx$$




Note: the area below the x -axis gives a negative value for its area. You must take that value as a positive value to determine the area between a curve and the x -axis. Sketching the graph will show what part of the function lies below the x -axis.

Area between two curves

Using definite integrals you can also find the areas enclosed between curves.

$$A_{\text{between}} = \int_a^b (g(x) - f(x)) dx$$



With $g(x)$ as the “top” function (furthest from the x -axis). For the area between curves, it does not matter what is above/below the x -axis.

5.1 Indefinite integral and boundary condition

Integration is essentially the opposite of derivation. The following equation shows how to integrate a function:

$$\int x^n dx = \frac{x^{n+1}}{n+1} + C, \quad n \neq -1$$

As you can see, every time you integrate the power on your variable will increase by 1 and you divide by the new power. This is opposite of what happens when you derive, then the power always decreases and you multiply by the original power.

Whenever you integrate you also **always add** $+C$ to this function. This accounts for any constant that may have been lost while deriving. As you may have noticed, whenever you do derivation any constants that were in the original function, $f(x)$, become 0 in the derivative function, $f'(x)$. In order to determine the value of C , you need to fill in a point that lies on the curve to set up an equation in which you can solve for C .

This is the same thing you need to do when finding the y -intercept, C , for a linear function – see Functions: Linear functions.

Standard integration

Let $f'(x) = 12x^2 - 2$
Given that $f(-1) = 1$, find $f(x)$.

1. Separate summed parts (optional)

$$\int 12x^2 - 2 dx = \int 12x^2 dx + \int -2 dx$$

2. Integrate

$$f(x) = \int 12x^2 dx + \int -2 dx = \frac{12}{3}x^3 - 2x + C$$

3. Fill in values of x and $f(x)$ to find C

$$\begin{aligned} \text{Since } f(-1) &= 1, \\ 4(-1)^3 - 2(-1) + C &= 1 \\ C &= 3 \end{aligned}$$

$$\text{So: } f(x) = 4x^3 - 2x + 3$$

5.1.1 Integration with an internal function

$$\int f(ax + b) dx \quad \text{integrate normally and multiply by } \frac{1}{\text{coefficient of } x}$$

Example.

Find the following integrals:

$$\int e^{3x-4} dx$$

Coefficient of $x = 3$, so

$$\int e^{3x-4} dx = \frac{1}{3}e^{3x-4} + C$$

$$\int \cos(5x-2) dx$$

Coefficient of $x = 5$, so

$$\int \cos(5x-2) dx = \frac{1}{5} \sin(5x-2) + C$$

5.1.2 Integration by substitution

$$\int f(g(x)) \cdot g'(x) dx$$

Usually these questions will be the most complicated-looking integrals you will have to solve. So if an integration question looks complicated, try to look for a function and its derivative inside the function you are looking to integrate; it is likely to be a question where you have to use the substitution method! Study the example to see how it's done.

Integrate by substitution	
Find $\int 3x^2 e^{x^3} dx$	
1. Identify the inside function u , this is the function whose derivative is also inside $f(x)$	$g(x) = u = x^3$
2. Find the derivative $u' = \frac{du}{dx}$	$\frac{du}{dx} = 3x^2$
3. Substitute u and $\frac{du}{dx}$ into the integral (this way dx cancels out)	$\int e^u \frac{du}{dx} dx = \int e^u du = e^u + C$
4. Substitute u back to get a function with x	$\int e^u + C = e^{x^3} + C$

5.2 Definite integral

If there are limit values indicated on your integral, you are looking to find a definite integral. This means that these values will be used to find a numeric answer rather than a function.

This is done in the following way, where the values for a and b are substituted as x -values into your indefinite integral:

$$\int_a^b f(x) dx = F(b) - F(a) \quad \text{where} \quad F = \int f(x) dx$$

Be careful, the order you substitute a and b into the indefinite integral is relevant for your answer:

$$\int_a^b f(x) dx = - \int_b^a f(x) dx$$

Solving definite integrals

Find $\int_3^7 12x^2 - 2 dx$, knowing that $F(x) = 4x^3 - 2x$

1. Find the indefinite integral (without $+C$)


$$\int_3^7 12x^2 - 2 dx = [4x^3 - 2x]_3^7$$

2. Fill in: $F(b) - F(a)$
(integral $x = b$) - (integral $x = a$)

$$= [4(7)^3 - 2(7)] - [4(3)^3 - 2(3)] \\ = 1256$$

5.2.1 Area


Area between a curve and the x-axis



By determining a definite integral for a function, you can find the area beneath the curve that is between the two x -values indicated as its limits.

DB 5.5

$$A_{\text{curve}} = \int_a^b f(x) dx$$




The area below the x -axis gives a negative value for its area. You must take that value as a positive value to determine the area between a curve and the x -axis. Sketching the graph will show what part of the function lies below the x -axis. So

$$A_{\text{curve}} = \int_a^b f(x) dx + \left| \int_b^c f(x) dx \right|$$

or

$$A_{\text{curve}} = \int_a^c |f(x)| dx$$

Area between two curves



Using definite integrals you can also find the areas enclosed between curves:

$$A_{\text{between}} = \int_a^b (g(x) - f(x)) dx$$

With $g(x)$ as the “top” function (furthest from the x -axis). For the area between curves, it does not matter what is above/below the x -axis.

Finding areas with definite integrals.


Let $y = x^3 - 4x^2 + 3x$
 Find the area from $x = 0$ to $x = 3$.

1. Find the x -intercepts: $f(x) = 0$

$$x^3 - 4x^2 + 3x = 0, \text{ using the GDC:}$$

$$x = 0 \text{ or } x = 1 \text{ or } x = 3$$

2. If any of the x -intercepts lie within the range, sketch the function to see which parts lie above and below the x -axis.



3. Setup integrals and integrate

$$\text{Left: } \int_0^1 x^3 - 4x^2 + 3x \, dx =$$

$$= \left[\frac{1}{4}x^4 - \frac{4}{3}x^3 + \frac{3}{2}x^2 \right]_0^1$$

$$= \left(\frac{1}{4} - \frac{4}{3} + \frac{3}{2} \right) - (0)$$

$$= \frac{5}{12}$$

$$\text{Right: } \int_1^3 x^3 - 4x^2 + 3x \, dx =$$

$$= \left[\frac{1}{4}x^4 - \frac{4}{3}x^3 + \frac{3}{2}x^2 \right]_1^3$$

$$= \left(\frac{1}{4}(3)^4 - \frac{4}{3}(3)^3 + \frac{3}{2}(3)^2 \right)$$

$$- \left(\frac{1}{4}(1)^4 - \frac{4}{3}(1)^3 + \frac{3}{2}(1)^2 \right)$$


$$= -\frac{8}{3}$$


4. Add up the areas (and remember areas are never negative!)


$$\frac{5}{12} + \frac{8}{3} = \frac{37}{12}$$


Alternatively, use the calculator to find areas


Calculate the area between $\int_0^3 x^3 - 4x^2 + 3x$ and the x -axis



Press 
 4: Calculus
 2: Numerical integral



Enter the boundaries and before putting the function.
 Press , choose 'abs('



Enter the function and place the variable (usually x) after d

In this case, the area is 3.083

PROBABILITY

Table of contents & cheatsheet

Definitions

Sample space the list of all possible outcomes.

Event the outcomes that meet the requirement.

Probability for event A , $P(A) = \frac{\text{Number of ways } A \text{ can happen}}{\text{all outcomes in the sample space}}$.

Dependent events two events are dependent if the outcome of event A affects the outcome of event B so that the probability is changed.

Independent events two events are independent if the fact that A occurs does not affect the probability of B occurring.

Conditional probability the probability of A , given that B has happened: $P(A|B) = \frac{P(A \cap B)}{P(B)}$.


6.2. Multiple events

66

Probabilities for successive events can be expressed through tree diagrams or a table of outcomes.

Table of outcomes		
	H	T
H	H,H	H,T
T	T,H	T,T

Tree diagram




In general, if you are dealing with a question that asks for the probability of:


- one event **and** another, you **multiply**
- one event **or** another, you **add**

6.1. Single events

64



Mutually exclusive




$$P(A \cup B) = P(A) + P(B)$$

$$P(A \cap B) = 0$$

Combined events


$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

$$P(A \cap B) = P(A) + P(B) - P(A \cup B)$$



$A \cup B$ (union)

$A \cap B$ (intersect)



If independent: $P(A \cap B) = P(A) \times P(B)$.

Compliment, A' where $P(A') = 1 - P(A)$

Exhaustive when everything in the sample space is contained in the events

6.3. Distributions

69

For a distribution by function the domain of X must be defined as $\sum P(X = x) = 1$.

Expected value $E(X) = \sum xP(X = x)$

Binomial distribution $X \sim B(n, p)$ used in situations with only 2 possible outcomes and lots of trials

$$P(X = x) = \binom{n}{r} p^r (1-p)^{n-r}$$

where $\binom{n}{r} = nCr = \frac{n!}{r!(n-r)!}$, n = number of trials,
 p = probability of success, r = number of success.


On calculator:

- Binompdf(n,p,r) $P(X = r)$
- Binomcdf(n,p,r) $P(x \leq r)$

Mean = np

Variance = npq

Normal distribution $X \sim N(\mu, \sigma^2)$




where μ = mean, σ = standard deviation

On calculator:

- normcdf(lower bound, upper bound, μ, σ)
- invnorm(area, μ, σ)

6.1 Single events (Venn diagrams)

Probability for single events can be visually expressed through Venn diagrams:



Sample space the list of all possible outcomes.


Event the outcomes that meet the requirement.

Probability for event A ,

$$P(A) = \frac{\text{Number of ways } A \text{ can happen}}{\text{all outcomes in the sample space}}$$

Here the shaded circle.

Imagine I have a fruit bowl containing 10 pieces of fruit: 6 apples and 4 bananas.



These events are also exhaustive as there is nothing outside of the events (nothing in the sample space).

I pick a piece of fruit. Below are some common situations with Venn diagrams.


Mutually exclusive

What is the probability of picking each fruit?

Events do not overlap

$$P(A \cup B) = P(A) + P(B)$$

$$P(A \cap B) = 0$$



$$P(A) = \frac{6 \text{ Apples}}{10 \text{ pieces of fruit}} = 0.6$$

$$P(B) = \frac{4 \text{ Bananas}}{10 \text{ pieces of fruit}} = 0.4$$

In independent events

$$P(A \cap B) = P(A) \times P(B).$$

It will often be stated in questions if events are independent.


Combined events

Example.

Of the apples 2 are red, 2 are green and 2 are yellow.
 What is the probability of picking a yellow apple?

The intersect is the area the events overlap.

$$P(A \cap B) = P(A) + P(B) - P(A \cup B)$$



A: apples

B: yellow fruit

$$P(\text{yellow apple}) = \frac{2 \text{ apples}}{10 \text{ pieces of fruit}} = 0.2$$


Example.

What is the probability of picking an apple or a yellow fruit?

The union is the area contain by both events.

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

When an event is exhaustive the probability of the union is 1.



A: apples

B: yellow fruit

Event is exhaustive so probability of union is 1.


Compliment

Example.

What is the probability of not picking a yellow fruit?

Everything that is not in the stated event.

$$P(A') = 1 - P(A)$$



A: apples

B: yellow fruit

$$P(B') = 1 - P(B) = 1 - 0.6 = 0.4$$


Conditional

Example.

What is the probability of picking an apple given I pick a yellow fruit?

The probability given that some condition is already in place.

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$




A: apples

B: yellow fruit

$$P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{0.2}{(0.2 + 0.4)} = \frac{1}{3}$$

You can think of this as using B as the sample space, or removing the non yellow apples from the fruit bowl before choosing.



6.2 Multiple events (tree Diagrams)



Dependent events two events are dependent if the outcome of event A affects the outcome of event B so that the probability is changed.

Independent events two events are independent if the fact that A occurs does not affect the probability of B occurring.

Conditional probability the probability of A, given that B has happened:

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

Questions involving dependent events will often involve elements that are drawn “without replacement”. Remember that the probabilities will be changing with each new set of branches.

Probabilities for successive events can be expressed through tree diagrams or a table of outcomes. Often at standard level you will deal with two successive events, but both methods can be used for more. In general, if you are dealing with a question that asks for the probability of:


- one event **and** another, you **multiply**
- one event **or** another, you **add**

Tree diagrams

Example.

Two disks are randomly drawn without replacement from a stack of 4 red and 5 blue disks. Draw a tree diagram for all outcomes.

The probability of drawing two red disks can be found by multiplying both probabilities of getting red $\left(\frac{4}{9} \times \frac{3}{8}\right)$.



The probabilities for each event should always add up to 1. The probabilities describing all the possible outcomes should also equal 1 (that is, the probabilities that we found by multiplying along the individual branches).

What is the probability to draw one red and one blue disk?

$P(\text{one red and one blue})$

$$\begin{aligned} & (P(R) \text{ and } P(B)) \quad \text{or} \quad (P(B) \text{ and } P(R)) \\ & (P(R) \times P(B)) \quad \quad \quad (P(B) \times P(R)) \\ & \frac{20}{72} \quad \quad \quad + \quad \quad \quad \frac{20}{72} \quad \quad \quad = \frac{40}{72} = \frac{5}{9} \end{aligned}$$

It is common for conditional probability questions to relate to previous answers.

What is the probability to draw at least one red disk?

$P(\text{at least one red})$

$$\begin{aligned} & P(R \text{ and } R) + P(B \text{ and } R) + P(R \text{ and } B) = 1 - P(B \text{ and } B) \\ & \frac{12}{72} \quad + \quad \frac{20}{72} \quad + \quad \frac{20}{72} \quad = 1 - \frac{20}{72} \quad = \frac{52}{72} = \frac{13}{18} \end{aligned}$$

What is the probability of picking a blue disc given that at least one red disk is picked?

$$P(\text{blue disk} \mid \text{at least one red disk}) = \frac{P(\text{one red disk and one blue disk})}{P(\text{at least one red disk})} = \frac{\frac{5}{9}}{\frac{13}{18}} = \frac{10}{13}$$

Tables of Outcomes

A table of outcomes has the possible outcomes for one event in the first row and the possible outcomes for another event in the first column. The table is then filled in with either the combination of these outcomes or the number of items (or probability) that fall into both events.

Example.

Table of outcomes for two flips of a fair coin

	H	T
H	H,H	H,T
T	T,H	T,T

Example.

Table of outcomes for three machines and the average number of defective and non-defective items they make.

	Defective	Non-defective
Machine I	6	120
Machine II	4	80
Machine III	10	150

6.3 Distributions



Probability distribution a list of each possible value and their respective probabilities.

Expected value $E(X) = \sum xP(X = x)$

DB 4.7

We can take any of the examples above and create a probability distribution from them. It is important to define the factor X for which the probability applies. Once tabulated we can use the distribution to find the expected value. It is best to think of this as the average value you would get if you repeated the action many times.

Probability distributions

A fair coin is tossed twice, X is the number of heads obtained.

1. Draw a sample space diagram

	H	T
H	H, H	H, T
T	T, H	T, T

2. Tabulate the probability distribution

x	0	1	2
$P(X = x)$	$\frac{1}{4}$	$\frac{1}{2}$	$\frac{1}{4}$

(The sum of $P(X = x)$ always equals 1)

3. Find the expected value of X : $E(X)$

$$E(X) = \sum xP(X = x)$$

$$= 0 \cdot \frac{1}{4} + 1 \cdot \frac{1}{2} + 2 \cdot \frac{1}{4} = 1$$

So if you toss a coin twice, you expect to get heads once.

6.3.1 Distribution by function

A probability distribution can also be given by a function.

The domain of X must be specified, as the sum of the probabilities must equal 1.

Probability distribution by function

$$P(X = x) = k \left(\frac{1}{3}\right)^{x-1} \text{ for } x = 1, 2, 3. \text{ Find constant } k.$$

1. Use the fact that $\sum P(X = x) = 1$ $k \left(\frac{1}{3}\right)^{1-1} + k \left(\frac{1}{3}\right)^{2-1} + k \left(\frac{1}{3}\right)^{3-1} = 1$
2. Simplify and solve for k $k + \frac{1}{3}k + \frac{1}{9}k = \frac{13}{9}k = 1. \text{ So, } k = \frac{9}{13}.$

6.3.2 Binomial distribution



Binomial distribution $X \sim B(n, p)$ used to find probabilities in situations with only 2 possible outcomes and lots of trials

$$P(X = x) = \binom{n}{r} p^r (1-p)^{n-r}$$

where $\binom{n}{r} = nCr = \frac{n!}{r!(n-r)!}$
 n = number of trials
 p = probability of success
 r = number of success

You can calculate values using binomial expansion from the algebra chapter. However binomial distribution questions are often found on calculator papers.

For questions asking for the probability of an exact outcome, $P(X = r)$, we use Binompdf on the GDC.

For questions asking for the probability of several consecutive values, $P(X \leq r)$, we use Binomcdf on the GDC.


Note that Binomcdf only calculates $P(X \leq r)$ or in words “at most the value of r ”. Therefore you must remember to transform the function depending on the wording in the questions:

- “Less than r ” $P(X < r) = P(X \leq r - 1)$
- “More than r ” $P(X > r) = 1 - P(X \leq r)$
- “At least r ” $P(X \geq r) = 1 - P(X \leq r - 1)$

6.3.3 Normal distribution

A normal distribution is one type of probability distribution which gives a bell-shape curve if all the values and their corresponding probabilities are plotted.

We can use normal distributions to find the probability of obtaining a certain value or a range of values. This can be found using the area under the curve; the area under the bell-curve between two x -values always corresponds to the probability for getting an x -value in this range. The total area under the normal distribution is always 1; this is because the total probability of getting any x -value adds up to 1 (or, in other words, you are 100% certain that your x -value will lie somewhere on the x -axis below the bell-curve).



Notation: $X \sim N(\mu, \sigma^2)$

Transform to standard N: $Z = \frac{x - \mu}{\sigma}$

On calculator:

normcdf (lower bound, upper bound, mean ($= \mu$), standard deviation ($= \sigma$))

invnorm (area, mean ($= \mu$), standard deviation ($= \sigma$))

Even though you will be using your GDC to find probabilities for normal distributions, it's always *very* useful to draw a diagram to indicate for yourself (and the examiner) what area or x -value you are looking for.





Finding a probability or percentage using normal distribution


The weights of pears are normally distributed with mean = 110 g and standard deviation = 8 g.
Find the percentage of pears that weigh between 100 g and 130 g

Sketch!

Indicate:

- The mean = 110 g
- Lower bound = 100 g
- Upper bound = 130 g
- And shade the area you are looking to find.



Press , choose
5: Probability
5: Distributions
2: Normal Cdf

Enter lower and upper boundaries, mean (μ) and standard deviation (σ).
For lower bound = $-\infty$,
set lower: -1E99
For upper bound = ∞ ,
set upper: 1E99





Press 


So 88.8% of the pears weigh between 100 g and 130 g.

Finding an x -value using normal distribution when the probability is given

The weights of pears are normally distributed with mean = 110 g and standard deviation = 8 g. 8% of the pears weigh more than m grams. Find m .

Sketch!



Press 
 5: Probability
 5: Distributions
 3: Inverse Normal

Enter probability (Area), mean (μ) and standard deviation (σ).
 The calculator assumes the area is to the left of the x -value you are looking for.
 So in this case:
 area = $1 - 0.08 = 0.92$

Press 

So $m = 121$, which means that 8% of the pears weigh more than 121 g.

Finding mean and standard deviation of a normal distribution


All nails longer than 2.4 cm (5.5%) and shorter than 1.8 cm (8%) are rejected. What is the mean and standard deviation length?

1. Write down equations

$$P(L < 1.8) = 0.08$$

$$P(L > 2.4) = 0.055$$

2. Draw a sketch!



3. Write standardized equation of the form $P(Z < \dots)$

$$P\left(Z < \frac{1.8 - \mu}{\sigma}\right) = 0.08$$

$$P\left(Z > \frac{2.4 - \mu}{\sigma}\right) = 0.055$$

$$P\left(Z < \frac{2.4 - \mu}{\sigma}\right) = 1 - 0.055 = 0.945$$

4. Use invnorm on calculator

$$\text{invnorm}(0.08, 0, 1) = -1.4051$$

$$\text{invnorm}(0.945, 0, 1) = 1.5982$$

5. Equate and solve

$$\begin{cases} \frac{1.8 - \mu}{\sigma} = -1.4051 \\ \frac{2.4 - \mu}{\sigma} = 1.5982 \\ \mu = 2.08 \\ \sigma = 0.200 \end{cases}$$

Here you are solving a pair of simultaneous equations. For a review see the Prior Knowledge section.

STATISTICS

Table of contents & cheatsheet

Definitions

Population the entire group from which statistical data is drawn (and which the statistics obtained represent).

Sample the observations actually selected from the population for a statistical test.

Random Sample a sample that is selected from the population with no bias or criteria; the observations are made at random.

Discrete finite or countable number of possible values (e.g. money, number of people)

Continuous infinite amount of increments (e.g. time, weight)

Note: continuous data can be presented as discrete data, e.g. if you round time to the nearest minute or weight to the nearest kilogram.

7.1. Descriptive statistics

76

For 1 variable data with frequency use 1-Var Stats on GDC.

Mean the average value

$$\bar{x} = \frac{\text{the sum of the data}}{\text{no. of data points}}$$

Mode the value that occurs most often

Median when the data set is ordered low to high and the number of data points is:

- odd, then the median is the middle value;
- even, then the median is the average of the two middle values.

Range largest x -value – smallest x -value

Variance $\sigma^2 = \frac{\sum f(x - \bar{x})^2}{n}$ *calculator only*

Standard deviation $\sigma = \sqrt{\text{variance}}$ *calculator only*

Grouped data data presented as an interval

Use the midpoint as the x -value in all calculations.

Q₁ first quartile = 25th percentile

Q₂ median = 50th percentile

Q₃ third quartile = 75th percentile

Q₃ – Q₁ interquartile range (IQR) = middle 50 percent


7.3. Statistical graphs

79


Frequency the number of times an event occurs in an experiment

Cumulative frequency the sum of the frequency for a particular class and the frequencies for all the classes below it


Histogram



Cumulative frequency



Box and whisker plot



7.4. Bivariate statistics

83

For analysis of data with two variables.

On GDC use LinReg ($ax+b$).

Regression Line ($r = ax + b$)

Can be used to interpolate unknown data.

Interpretation of r -values


The correlation between the two sets of data. Can be positive or negative.

r -value	correlation
$0.00 \leq r \leq 0.25$	very weak
$0.25 \leq r \leq 0.50$	weak
$0.50 \leq r \leq 0.75$	moderate
$0.75 \leq r \leq 1.00$	strong


Correlation does not mean causation.

Scatter diagrams


Perfect positive



No correlation



Weak negative



7.1 Descriptive statistics

The mean, mode and median, are all ways of measuring “averages”. Depending on the distribution of the data, the values for the mean, mode and median can differ slightly or a lot. Therefore, the mean, mode and median are all useful for understanding your data set.



Example data set: 6, 3, 6, 13, 7, 7 in a table:

x	3	6	7	13
frequency	1	2	2	1

Mean the average value, $\bar{x} = \frac{\text{the sum of the data}}{\text{no. of data points}} = \frac{\sum x}{n} = \frac{\sum fx}{\sum f}$
 e.g. $\bar{x} = \frac{3+6+6+7+7+13}{6} = \frac{1 \cdot 3 + 2 \cdot 6 + 2 \cdot 7 + 1 \cdot 13}{1+2+2+1} = 7$

Mode the value that occurs most often (highest frequency)
 e.g. The example data set has 2 modes: 6 and 7

Median the middle value when the data set is ordered low to high. Even number of values: the median is the average of the two middle values.
 Find for larger values as $n + \frac{1}{2}$.
 e.g. data set from low to high: 3, 6, 6, 7, 7, 13
 median = $\frac{6+7}{2} = 6.5$

Range largest x -value – smallest x -value
 e.g. range = $13 - 3 = 10$

Variance $\sigma^2 = \frac{\sum f(x - \bar{x})^2}{n}$ *calculator only*

Standard deviation $\sigma = \sqrt{\text{variance}}$ *calculator only*

Grouped data data presented as an interval, e.g. $10 < x \leq 20$ where:

- lower boundary = 10
- upper boundary = 20
- interval width = $20 - 10 = 10$
- mid-interval value (midpoint) = $\frac{20 + 10}{2} = 15$

Use the midpoint as the x -value in all calculations with grouped data.

Adding a constant to all the values in a data set or multiplying the entire data set by a constant influences the mean and standard deviation values in the following way:

Table 7.1: Adding or multiplying by a constant

	adding constant k	multiplying by k
mean	$\bar{x} + k$	$k \times \bar{x}$
standard deviation	σ	$k \times \sigma$



Q_1	the value for x so that 25% of all the data values are \leq to it first quartile	= 25 th percentile
Q_2	median	= 50 th percentile
Q_3	third quartile	= 75 th percentile
$Q_3 - Q_1$	interquartile range (IQR)	= middle 50 percent

Example.

Snow depth is measured in centimetres:
30, 75, 125, 55, 60, 75, 65, 65, 45, 120, 70, 110.

Find the range, the median, the lower quartile, the upper quartile and the interquartile range.

First always rearrange data into ascending order: 30, 45, 55, 60, 65, 65, 70, 75, 75, 110, 120, 125

- The range:

$$125 - 30 = 95 \text{ cm}$$

- The median: there are 12 values so the median is between the 6th and 7th value.

$$\frac{65 + 70}{2} = 67.5 \text{ cm}$$

- The lower quartile: there are 12 values so the lower quartile is between the 3rd and 4th value.

$$\frac{55 + 60}{2} = 57.5 \text{ cm}$$

- The upper quartile: there are 12 values so the lower quartile is between the 9th and 10th value.

$$\frac{75 + 110}{2} = 92.5 \text{ cm}$$

- The IQR

$$92.5 - 57.5 = 35 \text{ cm}$$

7.2 Sampling techniques

In order to do estimations on the whole population, it is required to create samples that can represent the population. There are different sampling techniques that achieve this goal, where all of them have different advantages, flaws and objectives. It is good to be able to know the difference between each one of them.



Convenience sampling the sampling done on the easiest available members of the population.

e.g. if you wanted to do a survey on students in your school, you would ask students you are personally familiar with.

Simple random sampling the sampling where each member of the population has an equal chance of being selected.

e.g. you could randomly put all students' names in a hat and randomly select sample members out of it.

Systematic sampling the sampling where the population is arranged or listed in a specific order and then elements from that list are selected at fixed intervals starting at a random point.

e.g. all student names could be written in an alphabetical list and every 10th student is chosen.

Stratified sampling the population is split into several smaller groups, known as "strata". Then a random sample is selected from each strata.

e.g. students could be split according to their age groups, so that randomly several students from each age group are taken into a sample.

Quota sampling similarly to stratified sampling the population is split into groups. However, sampling from each group is done in a non-random manner. E.g. sampling could be done in proportion to the size of each strata.

e.g. if the students are split into females and males with ratio 70% to 30%, then your sample should contain about 70% females and 30% males.


7.3 Statistical graphs




Frequency the number of times an event occurs in an experiment

Cumulative frequency the sum of the frequency for a particular class and the frequencies for all the classes below it


Age	17	18	19	20	21
No. of students	21	45	93	61	20
Cumulative freq.	21	66	159	220	240



A histogram is used to display the frequency for a specific condition. The frequencies (here: # of students) are displayed on the y -axis, and the different classes of the sample (here: age) are displayed on the x -axis. As such, the differences in frequency between the different classes assumed in the sample can easily be compared.



The cumulative frequency graph is used to display the development of the frequencies as the classes of the event increase. The graph is plotted by using the sum of all frequencies for a particular class, added to the frequencies for all the classes below it. The classes of the event (age) are displayed on the x -axis, and the frequency is displayed on the y -axis. The cumulative frequency graph always goes upwards, because the cumulative frequency increases as you include more classes.



Box and whisker plots neatly summarize the distribution of the data. It gives information about the range, the median and the quartiles of the data. The first and third quartiles are at the ends of the box, the median is indicated with a vertical line in the interior of the box, and the maximum and minimum points are at the ends of the whiskers.




Outliers will be any points lower than $Q_1 - 1.5 \times \text{IQR}$ and larger than $Q_3 + 1.5 \times \text{IQR}$ (IQR = interquartile range)

To identify the value of Q_1 , Q_2 and Q_3 , it is easiest to use the cumulative frequency graph. First, determine the percentage of the quartile in question. Second, divide the total cumulative frequency of the graph (i.e. the total sample size) by 100 and multiply by the corresponding percentage. Then, you will have found the frequency (y -value) at which 25% for Q_1 / 50% for Q_2 / 75% for Q_3 of the sample is represented. To find the x -value, find the corresponding x -value for the previously identified y -value.

Example.

Using the histogram, create a cumulative frequency graph and use it to construct a box and whisker diagram.




Write out the table for frequency and cumulative frequency.


Frequency of fish	20-30	30-40	40-50	50-60	60-70	70-80	80-90	90-100	100-110	110-120
Length of fish	2	3	5	7	11	5	6	9	1	1
Cumulative f.	2	5	10	17	28	33	39	48	49	50

Example.


Plot on cumulative frequency chart. Remember to use the midpoint of the date, e.g., 25 for 20–30.



Use graph to find Q_1 , Q_2 and Q_3 .






Plot box and whiskers.



GDC

Finding the mean, standard deviation and quartiles etc.




Find the descriptive statistics for the data used in the previous example, showing the ages of students.



Press off on , go to Lists and Spreadsheets. Enter x -values in L1 and, if applicable, frequencies in L2

Press menu , choose 4: Statistics
1: Stat Calculations

1: One-Variable Statistics



Enter Num of lists: 1.
Press OK


Enter names of columns you used to enter your x -list and frequency list and column where you would like the solutions to appear: a[], b[] and c[].
Press OK

mean = 19.06;
standard deviation = 1.06 etc.

7.4 Bivariate statistics

Bivariate statistics are about relationships between two different variables. You can plot your individual pairs of measurements as (x, y) coordinates on a scatter diagram. Analysing bivariate data allows you to assess the relationship between the two measured variables; we describe this relationship as correlation.

Scatter diagrams



Through statistical methods, we can predict a mathematical model that would best describe the relationship between the two measured variables; this is called regression. In your exam you will be expected to find linear regression models using your GDC.

7.4.1 Regression line

The regression line is a linear mathematical model describing the relationship between the two measured variables. This can be used to find an estimated value for points for which we do not have actual data. It is possible to have two different types of regression lines: y on x (equation $y = ax + b$), which can estimate y given value x , and x on y (equation $x = yc + d$), which can estimate x given value y . If the correlation between the data is perfect, then the two regression lines will be the same.

However one has to be careful when extrapolating (going further than the actual data points) as it is open to greater uncertainty. In general, it is safe to say that you should not use your regression line to estimate values outside the range of the data set you based it on.

7.4.2 Pearson's correlation coefficient ($-1 \leq r \leq 1$)

Besides simply estimating the correlation between two variables from a scatter diagram, you can calculate a value that will describe it in a standardised way. This value is referred to as Pearson's correlation coefficient (r).



$r = 0$ means no correlation.
 $r \pm 1$ means a perfect positive/negative correlation.

Interpretation of r -values:

r -value	$0 < r \leq 0.25$	$0.25 < r \leq 0.50$	$0.50 < r \leq 0.75$	$0.75 < r < 1$
correlation	very weak	weak	moderate	strong

Remember that correlation \neq causation.

Calculate r while finding the regression equation on your GDC. Make sure that STAT DIAGNOSTICS is turned ON (can be found in the MODE settings), otherwise the r -value will not appear.

When asked to "comment on" an r -value make sure to include both, whether the correlation is:


1. positive / negative
and
2. strong / moderate / weak / very weak

Bivariate-statistics type questions

The height of a plant was measured the first 8 weeks

Week x	0	1	2	3	4	5	6	7	8
Height (cm) y	23.5	25	26.5	27	28.5	31.5	34.5	36	37.5

1. Plot a scatter diagram



The line of best fit should pass through the mean point.


2. Use the mean point to draw a best fit line

$$\bar{x} = \frac{0 + 1 + 2 + \dots + 8}{9} = 3.56$$

$$\bar{y} = \frac{23.5 + 25 + \dots + 37.5}{9} = 30$$


3. Find the equation of the regression line
Using GDC

$$y = 1.83x + 22.7$$




Press **off** **on**, got to "Lists and Spreadsheets"

Enter x -values in one column (e.g A) and y -values in another column (e.g. B)




Press **menu**
4: Statistics
1: Stat Calculations
3: Linear Regression (mx+b)



Enter
X list: A [];
Y list: B[];
1st Result Column: C[]

Press **OK**



So, equation of regression line is $y = 1.83x + 22.7$ and Pearson's correlation (r -value) = 0.986

4. Comment on the result.

Pearson's correlation is $r = 0.986$, which is a strong positive correlation.

