



Mathematics: analysis and approaches

Higher level

Paper 2

9 May 2023

Zone A afternoon | Zone B morning | Zone C afternoon

Candidate session number

--	--	--	--	--	--	--	--	--	--

2 hours

Instructions to candidates

- Write your session number in the boxes above.
- Do not open this examination paper until instructed to do so.
- A graphic display calculator is required for this paper.
- Section A: answer all questions. Answers must be written within the answer boxes provided.
- Section B: answer all questions in the answer booklet provided. Fill in your session number on the front of the answer booklet, and attach it to this examination paper and your cover sheet using the tag provided.
- Unless otherwise stated in the question, all numerical answers should be given exactly or correct to three significant figures.
- A clean copy of the **mathematics: analysis and approaches formula booklet** is required for this paper.
- The maximum mark for this examination paper is **[110 marks]**.



Do **not** write solutions on this page.


Section B

Answer **all** questions in the answer booklet provided. Please start each question on a new page.

10. [Maximum mark: 13]

A weight suspended on a spring is pulled down and released, so that it moves up and down vertically.

The height, H metres, of the base of the weight above the ground can be modelled by the function $H(t) = a\cos(7.8t) + b$, for $a, b \in \mathbb{R}$ and $0 \leq t \leq 10$, where t is the time in seconds after the weight is released.



(a) Find the period of the function.


[2]

(This question continues on the following page)

Do not write solutions on this page.

(Question 10 continued)

The weight is released when its base is at a minimum height of 1 metre above the ground, and it reaches a maximum height of 1.8 metres above the ground. The graph of H is shown in the following diagram.



- (b) Find the value of
 - (i) a ;
 - (ii) b . [3]
 - (c) Find the number of times that the weight reaches its maximum height in the first five seconds of its motion. [2]
 - (d) Find the first time that the base of the weight reaches a height of 1.5 metres. [2]
- A camera is set to take a picture of the weight at a random time during the first five seconds of its motion.
- (e) Find the probability that the height of the base of the weight is greater than 1.5 metres at the time the picture is taken. [4]

Do **not** write solutions on this page.

11. [Maximum mark: 20]

A game of chance involves drawing **two** balls at random out of a box without replacement. The box initially contains r red balls and y yellow balls.

Let $P(YY)$ represent the probability of drawing two yellow balls from the box without replacement.

Consider a version of this game where it is known that $P(YY) = \frac{1}{3}$.

(a) Show that $2y^2 - 2(r+1)y + r - r^2 = 0$. [4]

(b) By solving the equation in part (a), show that $y = \frac{(r+1) + \sqrt{3r^2+1}}{2}$. [4]

(c) Find two pairs of values for r and y that satisfy the condition $P(YY) = \frac{1}{3}$. [4]

Now consider a similar game of chance that involves drawing **three** balls out of a box without replacement. The box initially contains 10 red balls and y yellow balls.

Let $P(YYY)$ represent the probability of drawing three yellow balls from the box without replacement.

(d) Find an expression for $P(YYY)$ in terms of y . [3]

A yellow ball is added so that the box now contains 10 red balls and $(y+1)$ yellow balls. The probability of drawing three yellow balls from the box without replacement is now twice the probability expressed in part (d).

(e) Find the initial number of yellow balls in the box. [5]

Do **not** write solutions on this page.

12. [Maximum mark: 21]

Consider the differential equation $\frac{dy}{dx} = \frac{x^2 + 3y^2}{xy}$, where $x > 0, y > 0$.

It is given that $y = 2$ when $x = 1$.

- (a) Use Euler's method with step length 0.1 to find an approximate value of y when $x = 1.1$. [2]
- (b) By solving the differential equation, show that $y = x\sqrt{\frac{9x^4 - 1}{2}}$. [8]
- (c) Find the value of y when $x = 1.1$. [1]
- (d) With reference to the concavity of the graph of $y = x\sqrt{\frac{9x^4 - 1}{2}}$ for $1 \leq x \leq 1.1$, explain why the value of y found in part (c) is greater than the approximate value of y found in part (a). [2]

The graph of $y = x\sqrt{\frac{9x^4 - 1}{2}}$ for $\frac{\sqrt{3}}{3} < x < 1$ has a point of inflexion at the point P.

- (e) By sketching the graph of an appropriate derivative of y , determine the x -coordinate of P. [2]

It can be shown that $\frac{d^2y}{dx^2} = \frac{-x^4 + x^2y^2 + 6y^4}{x^2y^3}$, where $x > 0, y > 0$.

- (f) Use this expression for $\frac{d^2y}{dx^2}$ to show that point P lies on the straight line $y = mx$ where the exact value of m is to be determined. [6]