

# Markscheme

**May 2024**

**Chemistry**

**Higher level**

**Paper 3**

© International Baccalaureate Organization 2024

All rights reserved. No part of this product may be reproduced in any form or by any electronic or mechanical means, including information storage and retrieval systems, without the prior written permission from the IB. Additionally, the license tied with this product prohibits use of any selected files or extracts from this product. Use by third parties, including but not limited to publishers, private teachers, tutoring or study services, preparatory schools, vendors operating curriculum mapping services or teacher resource digital platforms and app developers, whether fee-covered or not, is prohibited and is a criminal offense.

More information on how to request written permission in the form of a license can be obtained from <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

© Organisation du Baccalauréat International 2024

Tous droits réservés. Aucune partie de ce produit ne peut être reproduite sous quelque forme ni par quelque moyen que ce soit, électronique ou mécanique, y compris des systèmes de stockage et de récupération d'informations, sans l'autorisation écrite préalable de l'IB. De plus, la licence associée à ce produit interdit toute utilisation de tout fichier ou extrait sélectionné dans ce produit. L'utilisation par des tiers, y compris, sans toutefois s'y limiter, des éditeurs, des professeurs particuliers, des services de tutorat ou d'aide aux études, des établissements de préparation à l'enseignement supérieur, des fournisseurs de services de planification des programmes d'études, des gestionnaires de plateformes pédagogiques en ligne, et des développeurs d'applications, moyennant paiement ou non, est interdite et constitue une infraction pénale.

Pour plus d'informations sur la procédure à suivre pour obtenir une autorisation écrite sous la forme d'une licence, rendez-vous à l'adresse <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

© Organización del Bachillerato Internacional, 2024

Todos los derechos reservados. No se podrá reproducir ninguna parte de este producto de ninguna forma ni por ningún medio electrónico o mecánico, incluidos los sistemas de almacenamiento y recuperación de información, sin la previa autorización por escrito del IB. Además, la licencia vinculada a este producto prohíbe el uso de todo archivo o fragmento seleccionado de este producto. El uso por parte de terceros —lo que incluye, a título enunciativo, editoriales, profesores particulares, servicios de apoyo académico o ayuda para el estudio, colegios preparatorios, desarrolladores de aplicaciones y entidades que presten servicios de planificación curricular u ofrezcan recursos para docentes mediante plataformas digitales—, ya sea incluido en tasas o no, está prohibido y constituye un delito.

En este enlace encontrará más información sobre cómo solicitar una autorización por escrito en forma de licencia: <https://ibo.org/become-an-ib-school/ib-publishing/licensing/applying-for-a-license/>.

## Subject details: Chemistry higher level paper 3 Markscheme

Candidates are required to answer **ALL** questions in Section A [**15 marks**] and all questions from **ONE** option in Section B [**30 marks**].

Maximum total = [**45 marks**].

1. Each row in the “Question” column relates to the smallest subpart of the question.
2. The maximum mark for each question subpart is indicated in the “Total” column.
3. Each marking point in the “Answers” column is shown by means of a tick (✓) at the end of the marking point.
4. A question subpart may have more marking points than the total allows. This will be indicated by “**max**” written after the mark in the “Total” column. The related rubric, if necessary, will be outlined in the “Notes” column.
5. An alternative word is indicated in the “Answers” column by a slash (/). Either word can be accepted.
6. An alternative answer is indicated in the “Answers” column by “**OR**”. Either answer can be accepted.
7. An alternative markscheme is indicated in the “Answers” column under heading **ALTERNATIVE 1** etc. Either alternative can be accepted.
8. Words inside chevrons « » in the “Answers” column are not necessary to gain the mark.
9. Words that are underlined are essential for the mark.
10. The order of marking points does not have to be as in the “Answers” column, unless stated otherwise in the “Notes” column.
11. If the candidate’s answer has the same “meaning” or can be clearly interpreted as being of equivalent significance, detail and validity as that in the “Answers” column then award the mark. Where this point is considered to be particularly relevant in a question it is emphasized by **OWTTE** (or words to that effect) in the “Notes” column.
12. Remember that many candidates are writing in a second language. Effective communication is more important than grammatical accuracy.
13. Occasionally, a part of a question may require an answer that is required for subsequent marking points. If an error is made in the first marking point then it should be penalized. However, if the incorrect answer is used correctly in subsequent marking points then **follow through** marks should be awarded. When marking, indicate this by adding **ECF** (error carried forward) on the script.
14. Do **not** penalize candidates for errors in units or significant figures, **unless** it is specifically referred to in the “Notes” column.
15. If a question specifically asks for the name of a substance, do not award a mark for a correct formula unless directed otherwise in the “Notes” column. Similarly, if the formula is specifically asked for, do not award a mark for a correct name unless directed otherwise in the “Notes” column.
16. If a question asks for an equation for a reaction, a balanced symbol equation is usually expected, do not award a mark for a word equation or an unbalanced equation unless directed otherwise in the “Notes” column.
17. Ignore missing or incorrect state symbols in an equation unless directed otherwise in the “Notes” column.

Section A

Question			Answers	Notes	Total
1.	a		a leak «of gas from the system» ✓	Accept amount/volume/concentration $H_2O_2$ much lower. Do <b>not</b> accept answers that refer to $MnO_2$ .	1
1.	b	i	scientific claims must be falsifiable <b>OR</b> it should be possible to test a scientific prediction by an experiment ✓	Accept answers that imply this principle such as if hypothesis not supported underlying theory needs to be examined.	1
1.	b	ii	doubling mass will double surface area ✓ «doubling surface area will» double collision rate / frequency of collisions ✓	Accept increasing instead of doubling for both M1 and M2. Do <b>not</b> accept more/greater probability/chance of collisions.	2
1.	b	iii	temperature <b>OR</b> concentration «of hydrogen peroxide»/[ $H_2O_2$ ] ✓	Accept different specific surface area / particle size. Do <b>not</b> accept simply surface area.	1

(continued...)

(Question 1 continued)

Question			Answers	Notes	Total
1.	c	i	tangent drawn to curve at $t = 0$ s ✓ slope/gradient calculation ✓	Accept start of reaction for $t=0$ s for M1. Do not accept line instead of tangent for M1. Do <b>not</b> accept simply taking a derivative for M2.	2
1.	c	ii	yes <b>AND</b> rate is negligible when no catalyst is present <b>OR</b> no <b>AND</b> reaction still occurs even when no catalyst is present ✓	Accept yes <b>AND</b> rate is zero when no catalyst is present. Do <b>not</b> accept yes <b>AND</b> the line gives a better fit with the data.	1

Question	Answers	Notes	Total
2. a	<p><b>ALTERNATIVE 1</b></p> $n(\text{CuSO}_4) \llcorner = \frac{6.4 \text{ g}}{(63.55 + 32.07 + (4 \times 16.00)) \text{ g mol}^{-1}} = \frac{6.4 \text{ g}}{159.62 \text{ g mol}^{-1}} \gg$ <p>= 0.040 «mol»</p> <p><b>AND</b></p> $n(\text{H}_2\text{O}) \llcorner = \frac{(10.0 - 6.4) \text{ g}}{(2 \times 1.01) + 16.00 \text{ g mol}^{-1}} = \frac{3.6 \text{ g}}{18.02 \text{ g mol}^{-1}} \gg = 0.20 \text{ «mol» } \checkmark$ $n(\text{CuSO}_4) : n(\text{H}_2\text{O}) \llcorner = \frac{0.20 \text{ mol}}{0.04 \text{ mol}} \gg = 5.0 \checkmark$ <p><b>ALTERNATIVE 2</b></p> <p>% CuSO<sub>4</sub> in CuSO<sub>4</sub>•5H<sub>2</sub>O</p> $\llcorner = 100 \times \frac{63.55 + 32.07 + (4 \times 16.00) \text{ g mol}^{-1}}{63.55 + 32.07 + (9 \times 16.00) + (10 \times 1.01) \text{ g mol}^{-1}} = \frac{159.62 \text{ g mol}^{-1}}{249.72 \text{ g mol}^{-1}} \gg$ <p>= 63.92 ✓</p> <p>mass of CuSO<sub>4</sub> «= 10.0 g × <math>\frac{63.92}{100}</math> » = 6.392 g ✓</p> <p><b>ALTERNATIVE 3</b></p> <p>mass of H<sub>2</sub>O in CuSO<sub>4</sub>•5H<sub>2</sub>O</p> $\llcorner = 10 \text{ g} \times \frac{(5 \times 16.00) + (10 \times 1.01) \text{ g mol}^{-1}}{(63.55 + 32.07 + (9 \times 16.00) + (10 \times 1.01)) \text{ g mol}^{-1}} = \frac{900.1 \text{ g}}{249.72} \gg$ <p>= 3.6 g ✓</p> <p>mass of CuSO<sub>4</sub> in CuSO<sub>4</sub>•5H<sub>2</sub>O «= 10.0 g - 3.6 g» = 6.4 g</p> <p><b>OR</b></p> <p>comparison with graph shows mass lost is 3.6 g ✓</p>	<p>Accept other valid methods.</p>	<p>2</p>

(continued...)

(Question 2 continued)

Question			Answers	Notes	Total
2.	b	i	uncertainty in mass loss $\llcorner = 0.05 + 0.05 \llcorner = \llcorner \pm \llcorner 0.1 \text{ g } \checkmark$ percentage uncertainty in mass loss $\llcorner = \frac{0.1 \text{ g}}{(10.0 - 6.4)} \times 100 \llcorner = \llcorner \pm \llcorner 2.78 / 3 \% \checkmark$	Award <b>[2]</b> for correct final answer.  Accept 2.77.	2
2.	b	ii	absolute uncertainty in water of crystallization $\llcorner = 5 \times \frac{2.78}{100} \llcorner = \llcorner \pm \llcorner 0.14 \checkmark$	Accept $\llcorner \pm \llcorner 0.15$ if percentage uncertainty taken as 3%.  Do <b>not</b> award marks for answers with more or less significant figures.	1
2.	c		water molecules in different «chemical» environments <b>OR</b> water molecules have different type of bonding $\checkmark$  present in a 4:1 / 1:4 ratio $\checkmark$	Do <b>not</b> accept different kinds of water molecules.   Accept answers that clearly communicate the ratio.	2

**Section B**

**Option A — Materials**

Question			Answers	Notes	Total
3.	a	i	electronegativity difference $\Delta X \llcorner = 3.2 - 0.8 \llcorner = 2.4$ <b>AND</b> average electronegativity $\Sigma X \llcorner = \frac{1}{2} (3.2 + 0.8) \llcorner = 2.0 \checkmark$ falls in "ionic" region of bonding diagram $\checkmark$	Accept 75% ionic and 25% covalent for M2.  Do <b>not</b> accept ionic without reference to diagram.	2
3.	a	ii	electrostatic attraction <b>OR</b> attraction between oppositely charged ions $\checkmark$	Do <b>not</b> accept ionic.	1
3.	a	iii	«small» displacement brings ions of same charge close together <b>OR</b> «small» displacement results in repulsion «between same charged ions» $\checkmark$	Do <b>not</b> accept bonds are weak.	1
3.	b		all electrons are paired/have paired spins $\checkmark$		1
3.	c	i	«caesium» very reactive / very high in reactivity series <b>OR</b> cannot be reduced by chemical methods $\checkmark$	Do <b>not</b> accept simply caesium is reactive. Accept difficult to reduce «caesium ions». Accept Cs is higher on activity series than C.	1
3.	c	ii	$\text{Cs}^+(\text{l}) + \text{e}^- \rightarrow \text{Cs}(\text{l}) \checkmark$		1

(continued...)



(Question 3 continued)

Question			Answers	Notes	Total
3.	c	iii	amount of Cs $\llcorner = \frac{1.00 \text{ g}}{132.91 \text{ g mol}^{-1}} \llcorner = 0.00752 \llcorner \text{mol} \llcorner \checkmark$ charge $\llcorner = 0.00752 \text{ mol} \times 9.65 \times 10^4 \text{ C mol}^{-1} \llcorner = 726 \llcorner \text{C} \llcorner \checkmark$	Accept 0.007 for M1. Award [2] for correct final answer.	2
3.	d	i	<p><b>Similarity:</b>                      Any one of:                      increase rate of reaction <math>\checkmark</math>                      provide an alternative mechanism <math>\checkmark</math>                      reduce activation energy/<math>E_a</math> «for the reaction» <math>\checkmark</math>                      not consumed «in overall reaction» <math>\checkmark</math></p> <p><b>Difference:</b>                      Any one of:                      heterogeneous are in different phase to reactants <b>AND</b> homogeneous in same phase <math>\checkmark</math>                      heterogeneous are solids <b>AND</b> homogeneous are fluids/liquids/gases/ in solution <math>\checkmark</math>                      heterogeneous act on surface <b>AND</b> homogeneous act in fluid/medium <math>\checkmark</math>                      heterogeneous remain unchanged <b>AND</b> homogeneous consumed at one stage and regenerated at another <math>\checkmark</math>                      heterogeneous more easily removed «from product than homogenous» <math>\checkmark</math></p>	Accept state for phase.	2
3.	d	ii	«very» large surface area «to mass ratio» $\checkmark$		1
3.	d	iii	prevents oxidation $\checkmark$	Accept prevents reaction with air.	1


(continued...)

(Question 3 continued)

Question			Answers	Notes	Total
3.	d	iv	negative effect on/harmful to environment/human health ✓	Accept uncertain effect on human health. Do <b>not</b> award mark for poisonous/toxic.	1
3.	e	i	2 ✓		1
3.	e	ii	volume «= (614 pm) <sup>3</sup> » = 2.31 × 10 <sup>8</sup> «pm <sup>3</sup> » ✓	Accept 2.31 × 10 <sup>-22</sup> «cm <sup>3</sup> » <b>OR</b> 2.31 × 10 <sup>-28</sup> «m <sup>3</sup> ». Whilst units are not required, if given, they must correspond to the numerical value. Do <b>not</b> , for example, award the mark for 2.31 × 10 <sup>-22</sup> pm <sup>3</sup> .	1

(continued...)



(Question 3 continued)

Question			Answers	Notes	Total
3.	e	iii	<p><b>ALTERNATIVE 1:</b>                      volume of one mol « = <math>\frac{1}{2}(2.31 \times 10^{-22} \text{ cm}^3 \times 6.02 \times 10^{23})</math> »                      = 69.5 «cm<sup>3</sup>» ✓                      «density = <math>\frac{132.91 \text{ g mol}^{-1}}{69.5 \text{ cm}^3 \text{ mol}^{-1}}</math> =» 1.91 «g cm<sup>-3</sup>» ✓</p> <p><b>ALTERNATIVE 2:</b>                      volume of one mol of unit cells « = <math>2.31 \times 10^{-22} \text{ cm}^3 \times 6.02 \times 10^{23}</math> »                      = 139.1 «cm<sup>3</sup>» ✓                      «density = <math>\frac{2 \times 132.91 \text{ g mol}^{-1}}{139.1 \text{ cm}^3 \text{ mol}^{-1}}</math> =» 1.91 «g cm<sup>-3</sup>» ✓</p>	Award [2] for correct final answer.	2
3.	f		entropy change with a polydentate ligand more favourable/less negative «than change with many monodentate ligands» ✓	Accept chelate effect. Accept positive entropy change.	1
3.	g			Accept answers that include the C=O oxygens in the circle.	1

Question			Answers	Notes	Total
4.	a		increase flexibility ✓  reduce intermolecular forces between «polymer» chains ✓	Accept increase softness / reduce density / make less brittle for M1.  Accept increase distance between chains for M2. Accept allow chains to slide past each other more easily for M2.	2
4.	b		Class: «polychlorinated dibenzo» dioxin «s» ✓  Effect: hormone disrupting/disrupts metabolic processes <b>OR</b> cellular damage <b>OR</b> genetic damage <b>OR</b> carcinogenic ✓	Do <b>not</b> accept toxic (given in question stem) for M2.	2
4.	c	i	«facilitates sorting for» recycling ✓		1
4.	c	ii	A <b>AND</b> it has C=O/~1700 «cm <sup>-1</sup> » absorption ✓	Accept A <b>AND</b> B has C-Cl/~700 «cm <sup>-1</sup> » absorption.	1

(continued...)

(Question 4 continued)

Question			Answers	Notes	Total
4.	d		<p><b>Monomer A</b></p>  <p style="text-align: right;">✓</p> <p><b>Monomer B</b></p>  <p style="text-align: right;">✓</p>	<p>Accept any structures that have one -OH and one -COOH on each.</p> <p>Award <b>[1 max]</b> if one has two -OH and the other has two -COOH.</p> <p>Accept -COCl or -COOR instead of -COOH.</p>	2
4.	e	i	physical properties depend on molecular orientation «to a fixed axis in the material» ✓		1
4.	e	ii	dissolved <b>AND</b> phase transitions occur over range of concentrations <b>OR</b> only possible to vary concentration when dissolved <b>OR</b> pure liquids have «almost» fixed concentration ✓	Accept at any concentration for range of concentrations.	1

Option B — Biochemistry


Question			Answers	Notes	Total
5.	a		$6\text{CO}_2(\text{g}) + 6\text{H}_2\text{O}(\text{l}) \rightarrow \text{C}_6\text{H}_{12}\text{O}_6(\text{aq}) + 6\text{O}_2(\text{g}) \checkmark$	<i>Ignore light in the equation.</i>	1
5.	b	i	Type of reaction: condensation <b>AND</b> By-product: water $\checkmark$	<i>Accept addition–elimination for Type of reaction.</i>	1
5.	b	ii	insoluble «in water»/cytoplasm $\checkmark$	<i>Accept stored as solid grains.</i>	1
5.	b	iii	hydrolyze $\checkmark$	<i>Accept convert back to glucose. Accept break down starch «into smaller units/components/monomers».</i>	1

Question			Answers	Notes	Total
6.	a		corn oil <b>AND</b> highest content of polyunsaturated fatty acids <b>OR</b> corn oil <b>AND</b> highest number of C=C/double bonds «per molecule/triglyceride» $\checkmark$	<i>Do <b>not</b> accept least saturated/most unsaturated.</i>	1
6.	b		« $n(\text{I}_2) = 0.0330 \text{ dm}^3 \times 0.500 \text{ mol dm}^{-3} = \text{» } 0.0165 \text{ «mol» } \checkmark$ « $m(\text{I}_2) = 0.0165 \text{ mol} \times 253.8 \text{ g mol}^{-1} = \text{» } 4.19 \text{ «g» } \checkmark$ «iodine number = $\frac{4.19 \text{ g} \times 100}{5.00 \text{ g}}$ » = 83.8 $\checkmark$	<i>Award [3] for correct final answer. Award [2 max] for 41.9.</i>	3

(continued...)

(Question 6 continued)


Question		Answers	Notes	Total
6.	c	<p>higher proportion of «triglycerides of» saturated fatty acids  <b>OR</b>                      saturated fatty acids have greater surface area/higher electron density  <b>OR</b>                      molecules of «triglycerides of» saturated fatty acids are packed more closely/have a linear structure ✓</p> <p>stronger London/dispersion/instantaneous induced dipole-induced dipole forces «between linear chains» ✓</p>	<p>For M1 do <b>not</b> accept butter is saturated without reference to proportions.</p> <p>For M2 accept stronger intermolecular forces.</p> <p>Accept LDF for London dispersion forces</p>	2

Question		Answers	Notes	Total
7.	a	 <p>correct identification of both bonds ✓</p>	Accept circle including H.	1
7.	b	glycine/gly <b>AND</b> glutamic acid/glu <b>AND</b> cysteine/cys ✓	Accept structures. Accept in any order.	1

(continued...)

(Question 7 continued)

Question			Answers	Notes	Total
7.	c		glycine/gly ✓	Accept structure.	1
7.	d		glutamic acid/glu ✓	Accept structure. Apply ECF from 7(b).	1

Question			Answers	Notes	Total
8.	a		 <p>typical curve as shown ✓</p>	<p>Accept any curve with a single maximum.</p> <p>Ignore other annotations, such as temperature values on the x-axis.</p> <p>Do <b>not</b> penalize if curve does not touch the x-axis at high temperature.</p>	1

(continued...)



(Question 8 continued)


Question		Answers	Notes	Total
8.	b	partial replacement of «non-/scarcely biodegradable» detergent by biodegradable enzymes <b>OR</b> save energy as require lower temperature <b>OR</b> biodegradable ✓	Do <b>not</b> accept references to eutrophication, phosphates, nitrates, etc.  Accept break down naturally.	1
8.	c	Award <b>[1]</b> for any of: «treatment of» oil spills ✓ «treatment of harmful» effluents from sewage/paper mills/leather processing/textile industry/food industry ✓	Accept other reasonable examples.  Do <b>not</b> accept plastics.	1

Question		Answers	Notes	Total
9.	a	yes <b>AND</b> third spot «from bottom» $R_f = 0.36$ ✓	Accept values in the range of 0.35–0.37 for $R_f$ of third spot. Accept answers with YES implied, as in the third spot from the bottom has an $R_f$ of 0.36. Accept alternative working such as determination of $d$ using $R_f$ and $D$ , where $D = 64$ mm, to identify the third spot as lutein.	1

(continued...)


(Question 9 continued)

Question			Answers	Notes	Total
9.	b		«highly» conjugated bonds/ delocalized bonds <b>OR</b> «many» alternating single <b>AND</b> double/multiple bonds ✓ «intense» absorption «bands» in visible region ✓		2

Question	Answers	Notes	Total
<p><b>10</b></p> <p><b>a</b></p>	 <p><math>K_m</math>: 0.55 «<math>\times 10^{-3} \text{ mol dm}^{-3}</math>» ✓</p> <p><math>V_{max}</math>: 51 «<math>\times 10^{-3} \text{ mol dm}^{-3} \text{ min}^{-1}</math>» ✓</p>	<p>For M1, accept values in the range of 0.45–0.60 «<math>\times 10^{-3} \text{ mol dm}^{-3}</math>».</p> <p>Award [1 max] if both are correctly labelled on the graph without giving the values.</p>	<p><b>2</b></p>

(continued...)

(Question 10 continued)

Question		Answers	Notes	Total
10.	b	 <p>lower initial gradient <b>AND</b> tending towards same <math>V_{max}</math> at higher [substrate] ✓</p>	<p>Accept graph finishing at <math>V_{max}</math>.</p>	1

(continued...)

(Question 10 continued)

Question			Answers	Notes	Total
10.	c		Any two of: $K_m$ higher <b>OR</b> compete for the active site ✓ higher concentration of substrate/arachidonic acid to reach same $V_{max}$ ✓ slower production of prostaglandin «precursor» ✓		2 max

Question			Answers	Notes	Total
11.			«many» OH/hydroxyl «groups» ✓ can H-bond to water ✓	Accept alcohol/hydroxy, but <b>not</b> hydroxide for hydroxyl.	2

Question			Answers	Notes	Total
12.	a		H-bonds ✓ between «complementary» nitrogenous/purine/pyrimidine bases <b>OR</b> between amino and carbonyl groups ✓	Accept AT and CG for M2.	2
12.	b		TACCGTTCG ✓		1

Option C — Energy

Question		Answers	Notes	Total
13.	a	$\text{CH}_4(\text{g}) + 2\text{O}_2(\text{g}) \rightarrow \text{CO}_2(\text{g}) + 2\text{H}_2\text{O}(\text{l})$ ✓		1
13.	b	amount of $\text{CH}_4$ in 1.00 g clathrate «= $4 \times \frac{1 \text{ g}}{478.7 \text{ g mol}^{-1}}$ » = 0.008356 «mol» ✓ specific energy « = $0.008356 \text{ mol g}^{-1} \times 891 \text{ kJ mol}^{-1}$ » = 7.45 «kJ g <sup>-1</sup> » ✓	Award <b>[2]</b> for correct final answer.	2
13.	c	<i>Lower specific energy:</i> not all methane clathrate is methane <b>OR</b> part of methane clathrate does not burn ✓  <i>Higher energy density:</i> methane is gas <b>AND</b> methane clathrate is solid <b>OR</b> methane has much lower density «than methane clathrate» <b>OR</b> methane clathrate occupies a much smaller volume than «same mass of» methane ✓	<i>For M1 accept reference to water molecules in methane clathrate as indicating it is not all methane.</i>	2
13.	d	asymmetric «stretch» <b>AND</b> produces/changes dipole moment ✓		1

(continued...)

(Question 13 continued)

Question		Answers	Notes	Total
13.	e	methane more potent greenhouse gas/higher GWP than carbon dioxide «produced by burning it» ✓	<i>Accept answers that imply we could be harnessing the energy released by the combustion.</i>	1
13.	f	<p><i>Support:</i> overall increase in both the carbon dioxide concentration and the temperature «anomaly» ✓</p> <p><i>Refute:</i> CO<sub>2</sub> increases between ~1940 and ~1970 without corresponding increase in temperature <b>OR</b> global temperatures rose between ~1935 and ~1945 despite [CO<sub>2</sub>] remaining relatively constant <b>OR</b> increases in global temperature between ~1900 and ~1940 and between ~1970 and ~2010 are similar, but increase in CO<sub>2</sub> levels much higher in latter period ✓</p>	<p><i>For M1, accept there is a good/close correlation «in their shapes».</i></p> <p><i>For M2, accept the connection may just be coincidental «not causative».</i></p> <p><i>Do <b>not</b> accept comments about fluctuations.</i></p> <p><i>Accept years that are approximately the same as these listed.</i></p>	2

Question			Answers	Notes	Total
14.	a	i	methane produces less CO <sub>2</sub> per kJ of energy ✓  methane produces fewer pollutants/particulates/CO/VOCs ✓	<i>For M1 do <b>not</b> accept simply methane produces less CO<sub>2</sub>. For M2 accept gasoline undergoes incomplete combustion.</i>	2
14.	a	ii	greater energy density <b>OR</b> longer range «for same volume of fuel»/requires smaller fuel tank ✓	<i>Accept more concentrated source of energy. Accept greater distance between refuelling. Accept gasoline more readily available. Accept gasoline fuelled cars safer / have lower explosion/auto-ignition risk. Accept gasoline easier to handle / store / transport.</i>	1
14.	b	i	<b>A:</b> CH <sub>3</sub> OH <b>AND B:</b> CH <sub>3</sub> -O-CO-R <b>OR</b> <b>A:</b> C <sub>2</sub> H <sub>5</sub> OH <b>AND B:</b> C <sub>2</sub> H <sub>5</sub> -O-CO-R ✓	<i>Accept R-OH/R'-OH <b>AND</b> R-O-CO-R/R'-O-CO-R.  Do <b>not</b> accept names.</i>	1
14.	b	ii	low«er» viscosity ✓ «simple esters have» weaker London/dispersion forces «than vegetable oils» ✓	<i>Do <b>not</b> accept intermolecular forces or van der Waals' forces instead of London/dispersion forces.</i>	2



Question		Answers	Notes	Total
15.	a	<p><i>Similarity:</i> both convert mass into energy <b>OR</b> both increase the binding energy «per nucleon» ✓</p> <p><i>Difference:</i> fusion «two smaller/lighter» nuclei combine «to produce a larger/heavier one» <b>AND</b> fission «heavy» nucleus splits «into fragments/smaller/lighter nuclei» ✓</p>	<p><i>For similarity accept that mass is lost/mass of products is less than that of reactants.</i> <i>For similarity accept change of atomic number / new elements produced.</i></p> <p><i>Do <b>not</b> accept atoms/s instead of nucleus/i.</i></p>	2
15.	b	«dark» lines in spectrum of sunlight ✓	<p><i>Do <b>not</b> accept simply spectrometry/spectroscopy.</i> <i>Do <b>not</b> accept references to emission spectra.</i></p>	1
15.	c	«oxygen» free radical «species»/O <sub>2</sub> <sup>-</sup> /O <sub>2</sub> <sup>2-</sup> /HO <sub>2</sub> /HO ✓	<i>Do <b>not</b> penalize incorrect dots on radicals.</i>	1
15.	d	«nuclear reactor technology can be adapted for» nuclear weapons «production» ✓	<i>Do <b>not</b> accept geopolitical reasons.</i>	1

(continued...)

(Question 15 continued)

Question			Answers	Notes	Total
15.	e	i	low proportion/percentage/abundance of fissile/ <sup>235</sup> U isotope ✓	<i>Accept only one isotope/<sup>235</sup>U will undergo fission.</i>	1
15.	e	ii	diffusion/centrifuging requires a gas <b>OR</b> «only common» gaseous compound of uranium ✓		1
15.	e	iii	lighter molecules/ <sup>235</sup> UF <sub>6</sub> /containing <sup>235</sup> U diffuse more rapidly <b>OR</b> rate of diffusion inversely proportional to «square root of» molar mass ✓	<i>Accept U-238/heavier particles move to the outside wall of the centrifuge <b>OR</b> U-235/lighter particles stay in the middle.</i>  <i>Accept rate of diffusion decreases with increasing molar mass.</i>	1


Question		Answers	Notes	Total
16.	a	<p>light «energy» excites dye molecules  <b>OR</b>                      «excited» dye molecules oxidize/dye <math>\rightarrow</math> dye<sup>+</sup> + e<sup>-</sup>  <b>OR</b>                      electrons from dye pass into TiO<sub>2</sub> layer ✓</p> <p>oxidized dye molecules/dye<sup>+</sup> oxidize iodide/I<sup>-</sup> to tri-iodide/I<sub>3</sub><sup>-</sup>  <b>OR</b>  <math>2 \text{ dye}^+ + 3 \text{ I}^- \rightarrow 2 \text{ dye} + \text{ I}_3^-</math> ✓</p> <p>electrons flow from TiO<sub>2</sub> layer «through external circuit back» to counter electrode  <b>OR</b>                      tri-iodide/I<sub>3</sub><sup>-</sup> flows «through electrolyte» to counter electrode ✓</p> <p>electrons reduce tri-iodide/I<sub>3</sub><sup>-</sup> to iodide/I<sup>-</sup> «at counter electrode»  <math>\text{I}_3^- + 2\text{e}^- \rightarrow 3\text{I}^-</math> ✓</p>	<p><i>Accept iodine/I<sub>2</sub> for tri-iodide/I<sub>3</sub><sup>-</sup>.</i>  <i>Accept reversible arrows in equations.</i></p>	4

(continued...)

(Question 16 continued)

Question			Answers	Notes	Total
16.	b	i	<p>Positive electrode (cathode): lead(IV) oxide/PbO<sub>2</sub> ✓</p> <p>Negative electrode (anode): lead/Pb ✓</p>	<p>Do <b>not</b> accept lead(IV)/Pb<sup>4+</sup>.</p> <p>Accept correct reactants shown in half-equations.</p>	2
16.	b	ii	higher energy density/charge per unit mass ✓	<p>Do <b>not</b> accept lithium ion lighter/smaller without mentioning similar capacity.</p> <p>Do <b>not</b> accept references to toxicity.</p> <p>Accept does not involve a liquid/corrosive electrolyte.</p> <p>Accept lower internal resistance.</p>	1

Option D — Medicinal chemistry

Question			Answers	Notes	Total
17.	a		interfere with «bacteria» cell wall <b>OR</b> animal cells do not have a cell wall ✓	Do <b>not</b> accept transpeptidase without reference to cell wall.	1
17.	b	i	«bulky group/steric shield» prevents enzyme/penicillinase/beta-lactamase attacking $\beta$ -lactam ring ✓	Accept harder for penicillinase to access/reach $\beta$ -lactam ring.  Accept shields/increases stability of $\beta$ -lactam ring.	1
17.	b	ii		Circle must enclose the whole of the COOH group.	1

(continued...)

(Question 17 continued)

Question			Answers	Notes	Total
17.	c		<p>«low doses of» antibiotics reach the water/soil/animal waste</p> <p><b>OR</b></p> <p>«low doses of» antibiotics are present in the animals / food produced from animals ✓</p> <p>favours survival/spread of mutant/resistant bacteria ✓</p>	<p><i>Do <b>not</b> accept increased probability of mutation for M2.</i></p>	2

Question			Answers	Notes	Total
18.	a		<p><i>Any one of:</i></p> <p>addiction/dependency ✓</p> <p>decreased breathing-rate ✓</p> <p>decreased heart rates ✓</p> <p>constipation ✓</p> <p>reduced sex drive ✓</p> <p>loss of appetite ✓</p> <p>depression ✓</p>	<p><i>Do <b>not</b> accept vague responses such as suppression/decrease in brain activity.</i></p>	1
18.	b	i	<p>«phenolic» OH/hydroxyl replaced by OCH<sub>3</sub>/ether/methoxy group ✓</p>	<p><i>Accept methylated.</i></p> <p><i>Accept alcohol/hydroxy, but <b>not</b> hydroxide for hydroxyl.</i></p>	1

(continued...)

(Question 18 continued)


Question			Answers	Notes	Total
18.	b	ii	m/z = 299 ✓	Accept molecular ion = 300. Do <b>not</b> accept simply RMM = 300 Accept absence of m/z = 285. .	1
18.	b	iii	codeine has fewer OH/hydroxyl groups «than morphine» ✓ can cross the blood-brain barrier more easily ✓	Accept codeine is more lipid soluble / less polar «than morphine» for M1.	2

Question			Answers	Notes	Total
19.	a		anticoagulant/prevention of blood clots ✓	Accept blood thinner / reducing chance/incidence of strokes/heart attacks.	1
19.	b		synergistic effect <b>OR</b> increases anticoagulant effect ✓	Accept the effect of one reinforces the effect of the other. Accept increased «risk of» stomach/gastrointestinal bleeding.	1

Question		Answers	Notes	Total
20.	a	inhibits enzyme/H <sup>+</sup> K <sup>+</sup> -ATPase /gastric proton pump which secretes H <sup>+</sup> «ions into gastric juice» ✓	<i>Accept proton pump inhibitor/PPI.</i>	1
20.	b	$\text{pH} - \text{p}K_a = \log \frac{[\text{HCO}_3^-]}{[\text{H}_2\text{CO}_3]} = 1.12 \checkmark$ $[\text{HCO}_3^-] = \llcorner 13.18 \times 2.03 \times 10^{-3} = \llcorner 2.68 \times 10^{-2} \llcorner \text{«mol dm}^{-3}\llcorner \checkmark$	<i>Award [2] for correct final answer.</i>	2

Question		Answers	Notes	Total
21.	a	«drug» blocks/inhibits «viral» enzyme/neuraminidase/NA «activity» ✓ prevents virus from leaving/escaping host cells «thus cannot infect other cells» ✓		2
21.	b	zanamivir <b>AND</b> many OH/hydroxyl/NH/NH <sub>2</sub> groups «that can H-bond with water» ✓	<i>Accept alcohol/hydroxy, but <b>not</b> hydroxide for hydroxyl.</i>	1



Question		Answers	Notes	Total
22.	a	 <p style="text-align: right;">✓</p>	Both chiral centres must be identified for the mark.	1
22.	b	<p>chiral auxiliary binds to reactant blocking one reaction site «by steric hindrance»  <b>OR</b>                      chiral auxiliary creates stereochemical condition «necessary to follow a certain pathway»  <b>OR</b>                      stereochemical induction  <b>OR</b>                      existing chiral centre affects the configuration of new chiral centres ✓</p>	Accept «chemical synthesis» yields a racemic mixture/mixture of enantiomers/diastereomers.	1

Question			Answers	Notes	Total
23.	a		alpha emitters/radionuclides delivered by a carrier/drug/protein «directly to the cancer cells» ✓  alpha particles have low penetration <b>OR</b> «completely» absorbed within a short range ✓  high energy particles/highly ionising/high destructive power ✓	Accept lead-212 in M1.	3
23.	b	i	${}_{81}^{208}\text{Tl}$ ✓	Accept ${}^{208}\text{Tl}$ / thallium-208 / Tl-208.  Accept correct product shown in an equation.	1
23.	b	ii	<b>ALTERNATIVE 1:</b> $\lambda = \left\langle \frac{\ln 2}{61.0} \Rightarrow 1.136 \times 10^{-2} \text{ «min}^{-1}\text{»} \right\rangle$ ✓ $t = \left\langle \frac{\ln \frac{N_0}{N_t}}{\lambda} = \frac{2.303}{1.136 \times 10^{-2}} \Rightarrow 203 \text{ «min»} \right\rangle$ ✓  <b>ALTERNATIVE 2:</b> $t = \frac{\log \frac{N_t}{N_0}}{\log 0.5} \times t_{\frac{1}{2}}$ ✓ $t = \left\langle \frac{61}{0.301} \Rightarrow 203 \text{ «min»} \right\rangle$ ✓	Award [2] for the correct final answer.	2

Question		Answers				Notes	Total	
24.	a	<b>Compound</b>	<b>Number of signals</b>	<b>AND</b>	<b>Relative areas</b>	Award [1] for any two correct cells. Accept ratio of areas in any order. Do <b>not</b> apply ECF for ratios.	2	
		<i>Ibuprofen</i>	3		3:1:1			✓
		<i>Impurity X</i>	3		2:2:1			✓
24.	b	<p><b>B</b> <b>AND:</b></p> <p>absorbance at 3200–3600 «cm<sup>-1</sup>» <b>OR</b> absorbance due to O–H/hydroxyl in alcohol ✓</p> <p>absence of absorbance at 1700–1750 «cm<sup>-1</sup>» <b>OR</b> absence of «absorbance for» C=O/carbonyl «in carboxyl» ✓</p>				<p>“B” only necessary once.</p> <p>Award M2 for not A <b>AND</b> «absorbance for» C=O/carbonyl «in carboxyl group».</p> <p>Accept any absorbance value in the ranges given.</p>	2	