

# DP IB Maths: AA HL



Your notes

## 2.5 Reciprocal & Rational Functions

### Contents

- \* 2.5.1 Reciprocal & Rational Functions



Your notes

## 2.5.1 Reciprocal & Rational Functions

### Reciprocal Functions & Graphs

#### What is the reciprocal function?

- The **reciprocal function** is defined by  $f(x) = \frac{1}{x}$ ,  $x \neq 0$
- Its **domain** is the set of **all real values except 0**
- Its **range** is the set of **all real values except 0**
- The reciprocal function has a **self-inverse** nature
  - $f^{-1}(x) = f(x)$
  - $(f \circ f)(x) = x$

#### What are the key features of the reciprocal graph?

- The graph **does not have a y-intercept**
- The graph **does not have any roots**
- The graph has **two asymptotes**
  - A **horizontal** asymptote at the x-axis:  $y = 0$ 
    - This is the **limiting value** when the absolute value of x gets very large
  - A **vertical** asymptote at the y-axis:  $x = 0$ 
    - This is the value that causes the **denominator to be zero**
- The graph has **two axes of symmetry**
  - $y = x$
  - $y = -x$
- The graph **does not have any minimum or maximum points**



Your notes

## Linear Rational Functions & Graphs

### What is a rational function with linear terms?

- A **(linear) rational function** is of the form  $f(x) = \frac{ax + b}{cx + d}$ ,  $x \neq -\frac{d}{c}$
- Its **domain** is the set of **all real values except**  $-\frac{d}{c}$
- Its **range** is the set of **all real values except**  $\frac{a}{c}$
- The **reciprocal function** is a **special case** of a rational function

### What are the key features of linear rational graphs?

- The graph has a **y-intercept** at  $\left(0, \frac{b}{d}\right)$  provided  $d \neq 0$
- The graph has **one root** at  $\left(-\frac{b}{a}, 0\right)$  provided  $a \neq 0$
- The graph has **two asymptotes**
  - A **horizontal asymptote**:  $y = \frac{a}{c}$ 
    - This is the **limiting value** when the absolute value of  $x$  gets very large
  - A **vertical asymptote**:  $x = -\frac{d}{c}$ 
    - This is the value that causes the **denominator to be zero**
- The graph **does not have any minimum or maximum points**
- If you are asked to **sketch or draw** a rational graph:
  - Give the **coordinates** of any **intercepts** with the axes
  - Give the **equations** of the **asymptotes**

#### Examiner Tip

- If you draw a horizontal line anywhere it should only intersect this type of graph once at most
- The only horizontal line that should not intersect the graph is the horizontal asymptote
  - This can be used to check your sketch in an exam



Your notes

### Worked example

The function  $f$  is defined by  $f(x) = \frac{10 - 5x}{x + 2}$  for  $x \neq -2$ .

- a) Write down the equation of
- the vertical asymptote of the graph of  $f$ ,
  - the horizontal asymptote of the graph of  $f$ .

(i) Vertical asymptote is when denominator equals zero

$$x + 2 = 0 \quad \boxed{x = -2}$$

(ii) Horizontal asymptote is limiting value as  $x$  gets large

$$\lim_{x \rightarrow \infty} \frac{10 - 5x}{x + 2} = \lim_{x \rightarrow \infty} \frac{-5x}{x} \quad \boxed{y = -5}$$

- b) Find the coordinates of the intercepts of the graph of  $f$  with the axes.

$y$ -intercept occurs when  $x = 0$


$$y = \frac{10 - 5(0)}{0 + 2} = 5 \quad \boxed{(0, 5)}$$

$x$ -intercept occurs when  $y = 0$

$$\frac{10 - 5x}{x + 2} = 0 \Rightarrow 10 - 5x = 0 \Rightarrow x = 2 \quad \boxed{(2, 0)}$$

- c) Sketch the graph of  $f$ .

Include asymptotes and intercepts



Your notes

## Quadratic Rational Functions & Graphs

### How do I sketch the graph of a rational function where the terms are not linear?

- A rational function can be written  $f(x) = \frac{g(x)}{h(x)}$ 
  - Where  $g$  and  $h$  are polynomials
- To find the **y-intercept** evaluate  $\frac{g(0)}{h(0)}$
- To find the **x-intercept(s)** solve  $g(x) = 0$
- To find the equations of the **vertical asymptote(s)** solve  $h(x) = 0$
- There will also be an **asymptote** determined by what  $f(x)$  tends to as  $x$  approaches infinity
  - In this course it will be either:
    - **Horizontal**
    - **Oblique (a slanted line)**
  - This can be found by writing  $g(x)$  in the form  $h(x)Q(x) + r(x)$ 
    - You can do this by **polynomial division** or **comparing coefficients**
  - The function then tends to the curve  $y = Q(x)$

### What are the key features of rational graphs: quadratic over linear?

- For the rational function of the form  $f(x) = \frac{ax^2 + bx + c}{dx + e}$
- The graph has a **y-intercept** at  $\left(0, \frac{c}{e}\right)$  provided  $e \neq 0$
- The graph can have **0, 1 or 2 roots**
  - They are the solutions to  $ax^2 + bx + c = 0$
- The graph has **one vertical asymptote**  $x = -\frac{e}{d}$
- The graph has an **oblique asymptote**  $y = px + q$ 
  - Which can be found by writing  $ax^2 + bx + c$  in the form  $(dx + e)(px + q) + r$ 
    - Where  $p, q, r$  are constants
    - This can be done by **polynomial division** or **comparing coefficients**




Your notes



Your notes

$$y = \frac{ax^2 + bx + c}{dx + e}$$




Copyright © Save My Exams. All Rights Reserved

### What are the key features of rational graphs: linear over quadratic?

- For the rational function of the form  $f(x) = \frac{ax + b}{cx^2 + dx + e}$
- The graph has a **y-intercept** at  $\left(0, \frac{b}{e}\right)$  provided  $e \neq 0$
- The graph has **one root** at  $x = -\frac{b}{a}$
- The graph has can have **0, 1 or 2 vertical asymptotes**
  - They are the solutions to  $cx^2 + dx + e = 0$
- The graph has a **horizontal asymptote**

$$y = \frac{ax + b}{cx^2 + dx + e}$$



Copyright © Save My Exams. All Rights Reserved

### Examiner Tip

- If you draw a horizontal line anywhere it should only intersect this type of graph twice at most
  - This idea can be used to check your graph or help you sketch it



Your notes





Your notes

### Worked example

The function  $f$  is defined by  $f(x) = \frac{2x^2 + 5x - 3}{x + 1}$  for  $x \neq -1$ .

- a) (i) Show that  $\frac{2x^2 + 5x - 3}{x + 1} = px + q + \frac{r}{x + 1}$  for constants  $p$ ,  $q$  and  $r$  which are to be found.
- (ii) Hence write down the equation of the oblique asymptote of the graph of  $f$ .

(i) Write  $2x^2 + 5x - 3$  as  $(x+1)(px+q) + r$

$$2x^2 + 5x - 3 = px^2 + qx + px + q + r$$

Compare coefficients

$$\begin{array}{ccc} x^2 & x & \text{constant} \\ 2 = p & 5 = q + p & -3 = q + r \end{array}$$

$$\therefore p = 2 \quad \therefore q = 3 \quad \therefore r = -6$$

$$\frac{2x^2 + 5x - 3}{x + 1} = \frac{(x+1)(2x+3) - 6}{x+1} = 2x + 3 - \frac{6}{x+1}$$

(ii)  $y = 2x + 3$

- b) Find the coordinates of the intercepts of the graph of  $f$  with the axes.

y-intercept occurs when  $x = 0$

$$y = \frac{2(0)^2 + 5(0) - 3}{(0) + 1} = -3 \quad (0, -3)$$

x-intercept occurs when  $y = 0$


$$\frac{2x^2 + 5x - 3}{x + 1} = 0 \Rightarrow 2x^2 + 5x - 3 = 0 \Rightarrow (2x - 1)(x + 3) \Rightarrow x = 0.5 \text{ or } x = -3$$

$$(0.5, 0) \text{ and } (-3, 0)$$

- c) Sketch the graph of  $f$ .

Vertical asymptote when denominator is zero  $x = -1$

Include asymptotes and intercepts



Your notes