

# DP IB Maths: AA SL



Your notes

## 2.3 Functions Toolkit

### Contents

- \* 2.3.1 Language of Functions
- \* 2.3.2 Composite & Inverse Functions
- \* 2.3.3 Graphing Functions




Your notes

## 2.3.1 Language of Functions

### Language of Functions

#### What is a mapping?

- A **mapping transforms** one set of values (**inputs**) into another set of values (**outputs**)
- Mappings can be:
  - **One-to-one**
    - Each input gets mapped to **exactly one unique** output
    - No two inputs are mapped to the same output
    - For example: A mapping that cubes the input
  - **Many-to-one**
    - Each input gets mapped to **exactly one** output
    - Multiple inputs can be mapped to the same output
    - For example: A mapping that squares the input
  - **One-to-many**
    - An input can be mapped to **more than one** output
    - No two inputs are mapped to the same output
    - For example: A mapping that gives the numbers which when squared equal the input
  - **Many-to-many**
    - An input can be mapped to **more than one** output
    - Multiple inputs can be mapped to the same output
    - For example: A mapping that gives the factors of the input



Copyright © Save My Exams. All Rights Reserved




#### What is a function?




Your notes

- A **function** is a mapping between two sets of numbers where **each input** gets mapped to **exactly one output**
  - The output does not need to be unique
- **One-to-one** and **many-to-one** mappings are functions
- A mapping is a function if its graph passes the **vertical line test**
  - Any **vertical line** will intersect with the graph **at most once**



MANY-TO-ONE MAPPINGS ARE FUNCTIONS



ONE-TO-ONE MAPPINGS ARE FUNCTIONS

Copyright © Save My Exams. All Rights Reserved



### What notation is used for functions?

- Functions are denoted using letters (such as  $f$ ,  $v$ ,  $g$ , etc)
  - A function is followed by a variable in a bracket
  - This shows the input for the function
  - The letter  $f$  is used most commonly for functions and will be used for the remainder of this revision note
- $f(x)$  represents an expression for the value of the function  $f$  when evaluated for the variable  $x$
- Function notation gets rid of the need for words which makes it **universal**
  - $f = 5$  when  $x = 2$  can simply be written as  $f(2) = 5$


### What are the domain and range of a function?

- The **domain** of a function is the set of values that are used as **inputs**
- A domain should be stated with a function
  - If a domain is not stated then it is assumed the domain is all the real values which would work as inputs for the function
  - Domains are expressed in terms of the input
    - $x \leq 2$
- The **range** of a function is the set of values that are given as **outputs**
  - The range depends on the domain
  - Ranges are expressed in terms of the output



Your notes

- $f(x) \geq 0$
- To graph a function we use the **inputs as the x-coordinates** and the **outputs as the y-coordinates**
  - $f(2) = 5$  corresponds to the coordinates (2, 5)
- Graphing the function can help you visualise the range
- Common sets of numbers have special symbols:
  - $\mathbb{R}$  represents all the real numbers that can be placed on a number line
    - $x \in \mathbb{R}$  means  $x$  is a real number
  - $\mathbb{Q}$  represents all the rational numbers  $\frac{a}{b}$  where  $a$  and  $b$  are integers and  $b \neq 0$
  - $\mathbb{Z}$  represents all the integers (positive, negative and zero)
    - $\mathbb{Z}^+$  represents positive integers
  - $\mathbb{N}$  represents the natural numbers (0,1,2,3...)



Copyright © Save My Exams. All Rights Reserved

### What are piecewise functions?

- **Piecewise functions** are defined by different functions depending on which interval the input is in
  - E.g.  $f(x) = \begin{cases} x + 1 & x \leq 5 \\ 2x - 4 & 5 < x < 10 \\ x^2 & 10 \leq x \leq 20 \end{cases}$
- The region for the individual functions **cannot overlap**
- To evaluate a piecewise function for a particular value  $x = k$ 
  - Find which interval includes  $k$
  - Substitute  $x = k$  into the corresponding function



Your notes

- The function **may or may not be continuous** at the ends of the intervals
  - In the example above the function is
    - continuous at  $x = 5$  as  $5 + 1 = 2(5) - 4$
    - not continuous at  $x = 10$  as  $2(10) - 4 \neq 10^2$

### Examiner Tip

- Questions may refer to "the largest possible domain"
  - This would usually be  $x \in \mathbb{R}$  unless  $\mathbb{N}$ ,  $\mathbb{Z}$  or  $\mathbb{Q}$  has already been stated
  - There are usually some exceptions
    - e.g. square roots;  $x \geq 0$  for a function involving  $\sqrt{x}$
    - e.g. reciprocal functions;  $x \neq 2$  for a function with denominator  $(x - 2)$

### Worked example

For the function  $f(x) = x^3 + 1$ ,  $2 \leq x \leq 10$ :

- a) write down the value of  $f(7)$ .

Substitute  $x = 7$

$$f(7) = 7^3 + 1$$

$$f(7) = 344$$

- b) find the range of  $f(x)$ .

Find the values of  $x^3 + 1$  when  $2 \leq x \leq 10$

$$2 \leq x \leq 10$$

$$8 \leq x^3 \leq 1000$$

$$9 \leq x^3 + 1 \leq 1001$$

$$9 \leq f(x) \leq 1001$$



Your notes

## 2.3.2 Composite & Inverse Functions

### Composite Functions

#### What is a composite function?

- A **composite function** is where a function is applied to another function
- A composite function can be denoted
  - $(f \circ g)(x)$
  - $fg(x)$
  - $f(g(x))$
- The order matters
  - $(f \circ g)(x)$  means:
    - First apply  $g$  to  $x$  to get  $g(x)$
    - Then apply  $f$  to the previous output to get  $f(g(x))$
    - Always start with the function **closest to the variable**
  - $(f \circ g)(x)$  is not usually equal to  $(g \circ f)(x)$

#### How do I find the domain and range of a composite function?

- The domain of  $f \circ g$  is the set of values of  $X$ ...
  - which are a **subset** of the **domain of  $g$**
  - which maps  $g$  to a value that is in the **domain of  $f$**
- The range of  $f \circ g$  is the set of values of  $X$ ...
  - which are a **subset** of the **range of  $f$**
  - found by **applying  $f$**  to the **range of  $g$**
- To find the **domain** and **range** of  $f \circ g$ 
  - First find the **range of  $g$**
  - **Restrict** these values to the values that are **within the domain** of  $f$ 
    - The **domain** is the set of values that **produce the restricted range** of  $g$
    - The **range** is the set of values that are **produced using the restricted range** of  $g$  as the domain for  $f$
- For example: let  $f(x) = 2x + 1$ ,  $-5 \leq x \leq 5$  and  $g(x) = \sqrt{x}$ ,  $1 \leq x \leq 49$ 
  - The **range of  $g$**  is  $1 \leq g(x) \leq 7$ 
    - **Restricting** this to fit the **domain of  $f$**  results in  $1 \leq g(x) \leq 5$
  - The **domain** of  $f \circ g$  is therefore  $1 \leq x \leq 25$ 
    - These are the values of  $x$  which map to  $1 \leq g(x) \leq 5$
  - The **range** of  $f \circ g$  is therefore  $3 \leq (f \circ g)(x) \leq 11$ 
    - These are the values which  $f$  maps  $1 \leq g(x) \leq 5$  to

### Examiner Tip

- Make sure you know what your GDC is capable of with regard to functions
  - You may be able to store individual functions and find composite functions and their values for particular inputs
  - You may be able to graph composite functions directly and so deduce their domain and range from the graph
- $ff(x)$  is not the same as  $[f(x)]^2$



Your notes



Your notes

### Worked example

Given  $f(x) = \sqrt{x+4}$  and  $g(x) = 3 + 2x$ :

a) Write down the value of  $(g \circ f)(12)$ .

First apply function closest to input

$$(g \circ f)(12) = g(f(12))$$

$$f(12) = \sqrt{12+4} = \sqrt{16} = 4$$

$$g(4) = 3 + 2(4) = 11$$

$$(g \circ f)(12) = 11$$

b) Write down an expression for  $(f \circ g)(x)$ .

First apply function closest to input

$$(f \circ g)(x) = f(g(x))$$

$$= f(3+2x)$$

$$= \sqrt{3+2x+4}$$

$$(f \circ g)(x) = \sqrt{7+2x}$$

c) Write down an expression for  $(g \circ g)(x)$ .

$$(g \circ g)(x) = g(g(x))$$

$$= g(3+2x)$$

$$= 3 + 2(3+2x)$$

$$= 3 + 6 + 4x$$

$$(g \circ g)(x) = 9 + 4x$$





Your notes




Your notes

## Inverse Functions

### What is an inverse function?

- Only **one-to-one** functions have inverses
- A function has an inverse if its graph passes the **horizontal line test**
  - Any **horizontal line** will intersect with the graph **at most once**
- The **identity function**  $\text{id}$  maps each value to itself
  - $\text{id}(x) = x$
- If  $f \circ g$  and  $g \circ f$  have the **same effect as the identity function** then  $f$  and  $g$  are **inverses**
- Given a function  $f(x)$  we denote the **inverse function** as  $f^{-1}(x)$
- An inverse function **reverses the effect** of a function
  - $f(2) = 5$  means  $f^{-1}(5) = 2$
- Inverse functions are used to solve equations
  - The solution of  $f(x) = 5$  is  $x = f^{-1}(5)$
- A composite function made of  $f$  and  $f^{-1}$  has the **same effect as the identity function**
  - $(f \circ f^{-1})(x) = (f^{-1} \circ f)(x) = x$



Copyright © Save My Exams. All Rights Reserved




### What are the connections between a function and its inverse function?

- The **domain of a function** becomes the **range of its inverse**
- The **range of a function** becomes the **domain of its inverse**
- The graph of  $y = f^{-1}(x)$  is a **reflection** of the graph  $y = f(x)$  in the line  $y = x$ 
  - Therefore solutions to  $f(x) = x$  or  $f^{-1}(x) = x$  will also be solutions to  $f(x) = f^{-1}(x)$ 
    - There could be other solutions to  $f(x) = f^{-1}(x)$  that don't lie on the line  $y = x$



Your notes



Copyright © Save My Exams. All Rights Reserved



### How do I find the inverse of a function?

- STEP 1: **Swap** the  $x$  and  $y$  in  $y = f(x)$ 
  - If  $y = f^{-1}(x)$  then  $x = f(y)$
- STEP 2: **Rearrange**  $x = f(y)$  to make  $y$  the subject
- Note this can be done in any order
  - Rearrange  $y = f(x)$  to make  $x$  the subject
  - Swap  $x$  and  $y$

#### Examiner Tip

- Remember that an inverse function is a reflection of the original function in the line  $y = x$ 
  - Use your GDC to plot the function and its inverse on the same graph to visually check this
- $f^{-1}(x)$  is not the same as  $\frac{1}{f(x)}$



Your notes

### Worked example

For the function  $f(x) = \frac{2x}{x-1}$ ,  $x > 1$ :

a) Find the inverse of  $f(x)$ .

Let  $y = f^{-1}(x)$  and rearrange  $x = f(y)$

$$x = \frac{2y}{y-1}$$

$$x(y-1) = 2y$$

$$xy - x = 2y$$

$$xy - 2y = x$$

$$y(x-2) = x$$

$$y = \frac{x}{x-2}$$


$$f^{-1}(x) = \frac{x}{x-2}$$

b) Find the domain of  $f^{-1}(x)$ .

Domain of  $f^{-1}$  is the range of  $f$

Sketch  $y = f(x)$  to see range

For  $x > 1$ ,  $f(x) > 2$



$$\text{Domain of } f^{-1} : x > 2$$

c) Find the value of  $k$  such that  $f(k) = 6$ .

Use inverse  $f(a) = b \Leftrightarrow a = f^{-1}(b)$

$$k = f^{-1}(6) = \frac{6}{6-2}$$

$$k = \frac{3}{2}$$



Your notes



Your notes

## 2.3.3 Graphing Functions

### Graphing Functions

#### How do I graph the function $y = f(x)$ ?

- A point  $(a, b)$  lies on the graph  $y = f(x)$  if  $f(a) = b$
- The **horizontal axis** is used for the **domain**
- The **vertical axis** is used for the **range**
- You will be able to graph some functions by hand
- For some functions you will need to use your GDC
- You might be asked to graph the **sum** or **difference** of two functions
  - Use your GDC to graph  $y = f(x) + g(x)$  or  $y = f(x) - g(x)$
  - Just type the functions into the graphing mode

#### What is the difference between “draw” and “sketch”?

- If asked to sketch you should:
  - Show the general shape
  - Label any key points such as the intersections with the axes
  - Label the axes
- If asked to draw you should:
  - Use a pencil and ruler
  - Draw to scale
  - Plot any points **accurately**
  - Join points with a straight line or smooth curve
  - Label any key points such as the intersections with the axes
  - Label the axes

#### How can my GDC help me sketch/draw a graph?

- You use your GDC to plot the graph
  - Check the scales on the graph to make sure you see the full shape
- Use your GDC to find any key points
- Use your GDC to check specific points to help you plot the graph

## Key Features of Graphs

### What are the key features of graphs?


- You should be familiar with the following key features and know how to use your GDC to find them
- Local minimums/maximiums
  - These are points where the graph has a minimum/maximum for a small region
  - They are also called **turning points**
    - This is where the graph changes its direction between upwards and downwards directions
  - A graph can have multiple local minimums/maximiums
  - A local minimum/maximum is not necessarily the minimum/maximum of the whole graph
    - This would be called the **global** minimum/maximum
  - For quadratic graphs the minimum/maximum is called the **vertex**
- Intercepts
  - $y$  - intercepts are where the graph crosses the  $y$ -axis
    - At these points  $x = 0$
  - $x$  - intercepts are where the graph crosses the  $x$ -axis
    - At these points  $y = 0$
    - These points are also called the **zeros of the function** or **roots of the equation**
- Symmetry
  - Some graphs have lines of symmetry
    - A quadratic will have a vertical line of symmetry
- Asymptotes
  - These are lines which the graph will get closer to but not cross
  - These can be horizontal or vertical
    - Exponential graphs have horizontal asymptotes
    - Graphs of variables which vary inversely can have vertical and horizontal asymptotes



Your notes



Your notes



Copyright © Save My Exams. All Rights Reserved



### Examiner Tip

- Most GDC makes/models will not plot/show asymptotes just from inputting a function
  - Add the asymptotes as additional graphs for your GDC to plot
  - You can then check the equations of your asymptotes visually
  - You may have to zoom in or change the viewing window options to confirm an asymptote
- Even if using your GDC to plot graphs and solve problems sketching them as part of your working is good exam technique
  - Label the key features of the graph and anything else relevant to the question on your sketch





Your notes

### Worked example

Two functions are defined by

$$f(x) = x^2 - 4x - 5 \text{ and } g(x) = 2 + \frac{1}{x+1}.$$

a) Draw the graph  $y = f(x)$ .


Draw means accurately

Use GDC to find vertex, roots and y-intercepts

Vertex =  $(2, -9)$

Roots =  $(-1, 0)$  and  $(5, 0)$

y-intercept =  $(0, -5)$



b) Sketch the graph  $y = g(x)$ .



Your notes


Sketch means rough but showing key points

Use GDC to find  $x$  and  $y$ -intercepts and asymptotes

$$x\text{-intercept} = \left(-\frac{3}{2}, 0\right)$$

$$y\text{-intercept} = (0, 3)$$

Asymptotes :  $x = -1$  and  $y = 2$






Your notes

## Intersecting Graphs

### How do I find where two graphs intersect?

- Plot both graphs on your GDC
- Use the intersect function to find the intersections
- Check if there is more than one point of intersection



- LINES INTERSECT AT (2,1)
- SOLVING  $2x - y = 3$  AND  $3x + y = 7$  SIMULTANEOUSLY IS  $x = 2, y = 1$

Copyright © Save My Exams. All Rights Reserved



### How can I use graphs to solve equations?

- One method to solve equations is to use graphs
- To solve  $f(x) = a$ 
  - Plot the two graphs  $y = f(x)$  and  $y = a$  on your GDC
  - Find the points of intersections
  - The **x-coordinates** are the **solutions** of the equation
- To solve  $f(x) = g(x)$

- Plot the two graphs  $y = f(x)$  and  $y = g(x)$  on your GDC
- Find the points of intersections
- The **x-coordinates** are the **solutions** of the equation
- Using graphs makes it easier to see **how many solutions** an equation will have

### Examiner Tip

- You can use graphs to solve equations
  - Questions will not necessarily ask for a drawing/sketch or make reference to graphs
  - Use your GDC to plot the equations and find the intersections between the graphs



Your notes



Your notes


**Worked example**

Two functions are defined by

$$f(x) = x^3 - x \text{ and } g(x) = \frac{4}{x}$$


a) Sketch the graph  $y = f(x)$ .

Use GDC to find max, min, intercepts



b) Write down the number of real solutions to the equation  $x^3 - x = 2$ .

Identify the number of intersections between  $y = x^3 - x$  and  $y = 2$




c) Find the coordinates of the points where  $y = f(x)$  and  $y = g(x)$  intersect.



Your notes

Use GDC to sketch both graphs



$(-1.60, -2.50)$  and  $(1.60, 2.50)$

d)

Write down the solutions to the equation  $x^3 - x = \frac{4}{x}$ .

Solutions to  $x^3 - x = \frac{4}{x}$  are the  $x$  coordinates of the points of intersection.

$x = -1.60$  and  $x = 1.60$