

# DP IB Maths: AA SL



Your notes

## 5.5 Optimisation

### Contents

- \* 5.5.1 Modelling with Differentiation



Your notes

## 5.5.1 Modelling with Differentiation

### Modelling with Differentiation

#### What can be modelled with differentiation?

- Recall that **differentiation** is about the **rate of change** of a function and provides a way of finding **minimum** and **maximum** values of a function
- Anything that involves **maximising** or **minimising** a quantity can be modelled using differentiation; for example
  - minimising the cost of raw materials in manufacturing a product
  - the maximum height a football could reach when kicked
- These are called **optimisation** problems

#### What modelling assumptions are used in optimisation problems?

- The quantity being optimised needs to be dependent on a **single** variable
  - If other variables are initially involved, **constraints** or **assumptions** about them will need to be made; for example
    - minimising the cost of the **main** raw material – timber in manufacturing furniture say
    - the cost of screws, glue, varnish, etc can be fixed or considered **negligible**
- Other **modelling assumptions** may have to be made too; for example
  - ignoring air resistance and wind when modelling the path of a kicked football

#### How do I solve optimisation problems?

- In **optimisation** problems, letters other than **x**, **y** and **f** are often used including capital letters
  - **V** is often used for volume, **S** for surface area
  - **r** for radius if a circle, cylinder or sphere is involved
- **Derivatives** can still be found but be clear about which letter is representing the independent (**x**) variable and which letter is representing the dependent (**y**) variable
  - A GDC may always use **x** and **y** but ensure you use the correct letters throughout your working and final answer
- Problems often start by **linking** two connected quantities together – for example **volume** and **surface area**
  - Where more than one variable is involved, **constraints** will be given such that the quantity of interest can be rewritten in terms of **one** variable
- Once the quantity of interest is written as a function of a single variable, **differentiation** can be used to **maximise** or **minimise** the quantity as required

#### STEP 1

Rewrite the quantity to be optimised in terms of a single variable, using any constraints given in the question

#### STEP 2

Differentiate and solve the derivative equal to zero to find the "x"-coordinate(s) of any stationary points

### STEP 3

If there is more than one stationary point, or the requirement to justify the nature of the stationary point, differentiate again

### STEP 4

Use the second derivative to determine the nature of each stationary point and select the maximum or minimum point as necessary

### STEP 5

Interpret the answer in the context of the question

### Examiner Tip

- The first part of rewriting a quantity as a single variable is often a "show that" question – this means you may still be able to access later parts of the question even if you can't do this bit
- Even when an algebraic solution is required you can still use your GDC to check answers and help you get an idea of what you are aiming for




Your notes



Your notes

### Worked example

A large allotment bed is being designed as a rectangle with a semicircle on each end, as shown in the diagram below.



The total area of the bed is to be  $100\pi \text{ m}^2$ .

- a) Show that the perimeter of the bed is given by the formula

$$P = \pi \left( r + \frac{100}{r} \right)$$





Your notes

STEP 1: Rewrite  $P$  as powers of  $r$

$$P = \pi(r + 100r^{-1})$$

STEP 2:  $\frac{dP}{dr} = \pi(1 - 100r^{-2})$

$$\therefore \frac{dP}{dr} = \pi \left( 1 - \frac{100}{r^2} \right)$$

c) Find the value of  $r$  that minimises the perimeter.

STEP 2:  $\pi \left( 1 - \frac{100}{r^2} \right) = 0$

$$r^2 - 100 = 0$$

$$r = 10 \quad (\text{reject } -10 \text{ as } r \text{ is a length})$$

This is the only stationary point so we can assume it is minimal.

$$\therefore r = 10 \text{ m minimises the perimeter}$$

d) Hence find the minimum perimeter.



Your notes

STEP 5: Interpret answer in context

Minimum perimeter is when  $r=10$

$$\therefore P = \pi \left( 10 + \frac{100}{10} \right) = 20\pi$$

Minimum perimeter is  $20\pi$  m

Use your GDC to check

e) Justify that this is the minimum perimeter.

STEP 4: Use second derivative

$$\frac{d^2P}{dr^2} = \pi (200r^{-3})$$

$$\text{at } r=10, \frac{d^2P}{dr^2} = \frac{\pi}{5} > 0 \therefore \text{minimum}$$

$\therefore 20\pi$  is the minimum perimeter